


One gallon
Red
3.785L

quseptic

**GENERAL DISINFECTANT
AND DEODORANT**



Directions: Quseptic should be diluted for all uses. Thoroughly mix the required amounts of Quseptic and water before use. In all uses, the diluted disinfectant solution should be in contact with the object for at least 10 minutes.

General disinfectant for inanimate objects. (Kennels, operating tables and rooms, floors, refrigerators, lavatories.) Clean the object with a detergent and rinse thoroughly. Disinfect non-porous surfaces by applying a solution of 1 oz. (approximately 2 tablespoonfuls) of Quseptic in a gal. of water. For porous surfaces use 2 oz. of Quseptic per gal. of water.

As a sanitizer for utensils, glassware, bottles. Clean with a suitable detergent, rinse thoroughly, then dip in a solution of 1 oz. to 4 gal. water and follow with a potable water rinse.

As a sanitizer for dairy equipment. Clean milking machines, milk cans, etc., with detergent and rinse thoroughly. Then immerse in a solution of 1/4 oz. (approx. 2 teaspoonfuls) Quseptic in 1 gal. water. Follow with a potable water rinse. This same strength solution may be used in the teat cups while they are hanging in the racks. This diluted solution is also recommended as a sanitizing wash for wiping the flank and udder prior to milking.

For disinfecting rubber goods, towels, clothes. Clean the object with detergent, rinse thoroughly. Soak for 10 minutes in solution of 1 oz. to 1 gal. of water. Completely immerse fabric. Three gal. is sufficient for 10 lb. of dry cloth.

For disinfecting instruments. Clean with suitable detergent and rinse thoroughly. Immerse instruments for 10 minutes in solution containing 1 oz. to 1 gal. of water.

For deodorizing. Use 1/2 to 1 oz. Quseptic to 1 gal. water. Spray on kennels, runways, garbage cans and other places to aid in killing odors without masking.


Warning: May cause skin irritation or damage to the eyes. Do not get in eyes. Avoid contact with the skin. In case of contact, flush immediately with plenty of water. For eyes, get medical attention. Harmful if swallowed. Avoid contamination of food.

Soap reduces the germicidal efficiency of Quseptic, and care should be exercised to rinse an object thoroughly before it is placed in contact with various diluted solutions of Quseptic. Detergents such as trisodium phosphate, tetrasodium pyro-phosphate, soda ash, sodium sesquicarbonate, etc., are compatible.

Note: If it is so desired, a stock solution may be prepared by diluting one gallon Quseptic with three gal. water. This stock solution is then used the same as directed above except that four times more is required, i.e., 1 oz. is diluted to 1 gal. instead of 1/4 oz. to 1 gal.

Lot No.

Norden Laboratories, Lincoln, Nebraska 68501, U.S.A.
Downsview, Ontario Calgary, Alberta



One gallon
Red
3.785L


quseptic

GENERAL DISINFECTANT
AND DEODORANT

ACCEPTED

Aug. 16, 1972

UNIVERSITY OF MICHIGAN
FEDERAL BUREAU OF INVESTIGATION
RECEIVED
ANN ARBOR, MI 48106-15



quseptic

GENERAL DISINFECTANT
AND DEODORANT

Contains: Active ingredients:

n - (Alkyl [50% C ₁₄ , 40% C ₁₂ , 10% C ₁₆]) dimethyl benzyl ammonium chloride.....	10.0%
Ethyl alcohol	2.5%
Inert ingredients	87.5%

Phenol coefficients (A.O.A.C. method, 20°C):

Salmonella typhosa—46.
Staphylococcus aureus—70.

Note: This product fulfills the criteria of Appendix F of the Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of the U.S. Public Health Service in waters up to 450 ppm of hardness calculated as CaCO₃ when tested by the A.O.A.C. Germicidal and Detergent Sanitizers—Official Method.

WARNING: Keep Out of Reach of Children.

Rinse empty container thoroughly with water and discard it.

EPA Reg. No. 622-15

For Veterinary Use Only

Lot No.

Norden Laboratories, Lincoln, Nebr. 68501, U.S.A.

80-7850

 **NORDEN**
LABORATORIES