


linspray

(CONTAINS LINDANE)

One Gallon



Directions

Beef cattle and horses—Ticks, lice, horn fly, mosquitoes: Spray with Dilution No. 2. Repeat treatment as necessary.

Sheep—Ticks: Use Dilution No. 1 and spray along back and around ears, spraying against wool using high pressure.

Hogs—Mange (Sarcoptes scabiei) and lice: Use Dilution No. 2.

Sheep—Sheep scab: Spray thoroughly with high pressure, using Dilution No. 2.

As a dip—For control of listed pests: Use Dilution No. 2. Fill vat with water, then add the required amount of LINSPRAY (See Dilution No. 2) and stir thoroughly for 3 minutes or longer. If dipping is stopped longer than 30 minutes, agitate vat contents thoroughly before resuming dipping. Completely submerge animal's body once or twice and duck head at least twice. Do not dip young calves or over heated animals. Water animals before dipping to prevent drinking of solution. Treat calves only at the 0.03% level. Do not dip calves under 3 months of age.

As a Paint: Paint windows, doors, ceilings, and other favorable places where flies congregate. Dilute one part LINSPRAY with 9 parts water. Do not use in dairy barns.


Residual spray for dog kennels, pens, and manure piles for control of flies, roaches, fleas, mites, mosquitoes and ants: Spray thoroughly using Dilution No. 3.

DILUTION TABLE

Amount of Water to Use	Amount of LINSPRAY to Use		
	Dilution No. 1 0.03% Lindane	Dilution No. 2 0.06% Lindane	Dilution No. 3 0.10% Lindane
1 Gallon	2 Teaspoons	4 Teaspoons	6 Tablespoons
5 Gallons	1 Tablespoon	2 Tablespoons	1 Pint
50 Gallons	1 pint	2 Pints	1 1/2 Gallons

Lot No

Norden Laboratories, Lincoln, Nebraska 68501, U.S.A.



Indications: For use in spraying or dipping livestock for control of lice, ticks, fleas, mange, flies, mosquitoes; and for residual spray uses in pens, dog kennels and manure piles for control of flies, roaches, fleas, mites, mosquitoes and ants.

Directions

Beef cattle and horses—Ticks, lice, horn fly, mosquitoes: spray with Dilution No. 2. Repeat treatment as necessary.

Sheep—Ticks: Use Dilution No. 1 and spray along back and around ears, spraying against wool using high pressure.

Hogs—Mange (Sarcoptes scabiei) and lice: Use Dilution No. 2.

Sheep—Sheep scab: Spray thoroughly with high pressure, using Dilution No. 2.

As a dip—For control of listed pests: Use Dilution No. 2. Fill vat with water, then add the required amount of LINSPRAY (See Dilution No. 2) and stir thoroughly for 3 minutes or longer. If dipping is stopped longer than 30 minutes, agitate vat contents thoroughly before resuming dipping. Completely submerge animal's body once or twice and duck head at least twice. Do not dip young calves or over heated animals. Water animals before dipping to prevent drinking of solution. Treat calves only at the 0.03%

level. Do not dip calves under 3 months of age.

As a Paint: Paint windows, doors, ceilings and other favorable places where flies congregate. Dilute one part LINSPRAY with 9 parts water. Do not use in dairy barns.

Residual spray for dog kennels, pens, and manure piles for control of flies, roaches, fleas, mites, mosquitoes and ants: Spray thoroughly using Dilution No. 3.

DILUTION TABLE

Amount of Water to Use	Amount of LINSPRAY to Use		
	Dilution No. 1 0.03% Lindane	Dilution No. 2 0.06% Lindane	Dilution No. 3 0.10% Lindane
1 Gallon	2 Teaspoons	4 Teaspoons	6 Tablespoons
5 Gallons	1 Tablespoon	2 Tablespoons	1 Pint
50 Gallons	1 Pint	2 Pints	1 1/2 Gallons


Lot No

Norden Laboratories, Lincoln, Nebraska 68501, U.S.A.

One Pint

linspray

(Contains Lindane)



linspray

(CONTAINS LINDANE)

Indications: For use in spraying or dipping livestock for control of lice, ticks, fleas, mange, flies, mosquitoes; and for residual spray uses in pens, dog kennels and manure piles for control of flies, roaches, fleas, mites, mosquitoes and ants.

Active ingredients:
Gamma isomer of benzene hexachloride from lindane 12.0%
Aromatic petroleum derivative solvent..... 84.8%
Inert ingredients 3.2%

100.0%

CAUTION: Avoid inhalation. Avoid contact with the skin and eyes. May be absorbed through skin. Do not use on household pets or humans. In case of contact, wash immediately with soap and water. For eyes, get medical attention. Flush with water for 15 minutes.

Treat young animals very lightly and with caution. Avoid contamination of feed or foodstuffs. Cover feeding troughs before spraying. Do not apply with lime in fresh whitewash or with other highly alkaline materials. Do not apply to dairy animals. Do not dip animals within 60 days of slaughter. Do not spray within 30 days of slaughter. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather.

This product is toxic to fish and wildlife. Do not contaminate water supplies by cleaning of equipment or disposal of waste and containers. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

WARNING—FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.

EPA Reg. No. 622-31

For Veterinary Use Only

NORDEN
LABORATORIES

linspray

(CONTAINS LINDANE)

Active ingredients:
Gamma isomer of benzene hexachloride from lindane 12.0%
Aromatic petroleum derivative solvent..... 84.8%
Inert ingredients..... 3.2%

100.0%

CAUTION: Avoid inhalation. Avoid contact with the skin and eyes. May be absorbed through skin. Do not use on household pets or humans. In case of contact, wash immediately with soap and water. For eyes, get medical attention. Flush with water for 15 minutes.

Treat young animals very lightly and with caution. Avoid contamination of feed or foodstuffs. Cover feeding troughs before spraying. Do not apply with lime in fresh whitewash or with other highly alkaline materials. Do not apply to dairy animals. Do not dip animals within 60 days of slaughter. Do not spray within 30 days of slaughter. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather.

This product is toxic to fish and wildlife. Do not contaminate water supplies by cleaning of equipment or disposal of waste and containers. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

WARNING—FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.

EPA Reg. No. 622-31

For Veterinary Use Only

NORDEN
LABORATORIES