

AUG 16 1974

Label Copy  
6.5 gm.

622-38

6.5 gm.

MITOX OINTMENT

An aid in the treatment of bacterial infections of the external ear canal associated with ear mite infestation

Active Ingredients

Neomycin base (as the sulfate) .....	0.5%
Sulfacetamide .....	9.0%
Carbaryl (1-Naphthyl N-methylcarbamate)* .....	1.0%
Tetracaine hydrochloride .....	0.5%
Mineral oil .....	59.1%
Inert Ingredients .....	29.9%
	<u>100.0%</u>

\*Sevin

Store in cool place. See carton for directions.

NOTE: After using, retract plunger 1/16 inch to stop the flow of ointment.

CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED.

EPA Est. 622-NB-i

Lot No.

EPA Reg. No. 622-38

85-7048

For Veterinary Use Only

Norden Laboratories, Inc., Lincoln, Nebr. 68501, U.S.A.

N NORDEN  
LABORATORIES

Carton Copy  
12-6.5 gm. syringes

Front and Back Panel:

12-6.5 gm. syringes

MITOX OINTMENT

An aid in the treatment of bacterial infections of the  
external ear canal associated with ear mite infestation

CAUTION: Keep out of reach of children. Harmful if swallowed.

For Veterinary Use Only

**N** NORDEN  
LABORATORIES

Carton Copy  
12-6.5 gm. syringes

Left Panel:

12-6.5 gm. syringes  
Dispense-Pak

MITOX OINTMENT

Active Ingredients

Neomycin base (as the sulfate) .....	0.5%
Sulfacetamide .....	9.0%
Carbaryl (1-Naphthyl N-methylcarbamate)* .....	1.0%
Tetracaine hydrochloride .....	0.5%
Mineral Oil .....	59.1%
Inert Ingredients .....	29.9%
	<hr/>
	100.0%

\*Sevin

Lot No.

Carton Copy  
12-6.5 gm. syringes

Right Panel:

12-6.5 gm. syringes  
Dispense-Pak

#### MITOX OINTMENT

**Directions:** If severe accumulations of ceruminous material are present, clean ear before treatment with a ceruminolytic agent. Follow this cleaning with treatment with Mitox, as directed. Fill lower section with Mitox and massage gently to distribute evenly. For infections, repeat as needed. For ear mites, continue treatments at 10 day intervals until controlled.

**NOTE:** After using, retract plunger 1/16 inch to stop the flow of ointment before replacing cap.

EPA Reg. No. 622-38

EPA Est. 622-NB-1

Norden Laboratories, Lincoln, Nebraska 68501, U.S.A.

Carton Copy  
6.5 gm.

Dispensing Panel:

MITOX OINTMENT

An aid in the treatment of bacterial infections of the  
external ear canal associated with ear mite infestation

MITOX OINTMENT FOR DOGS AND CATS 6.5 gm.

**Active Ingredients**

Neomycin base (as the sulfate) .....	0.5%
Sulfacetamide .....	9.0%
Carbaryl (1-Naphthyl N-methylcarbamate)* .....	1.0%
Tetracaine hydrochloride .....	0.5%
Mineral oil .....	59.1%
<b>Inert Ingredients</b> .....	29.9%
	100.0%

\*Sevin

**Directions:** If severe accumulations of ceruminous material are present, clean ear before treatment with a ceruminolytic agent. Follow this cleaning with treatment with Mitox, as directed. Fill lower section with Mitox and massage gently to distribute evenly. For infections, repeat as needed. For ear mites, continue treatments at 10 day intervals until controlled.

**CAUTION:** Keep out of reach of children. Harmful if swallowed.

**NOTE:** After using, retract plunger 1/16 inch to stop the flow of ointment

For Veterinary Use Only

EPA Reg. No. 672-38

EPA Est. 672-NB-1

Norden Laboratories, Lincoln, Nebr. 68501, U.S.A.

|| NORDEN  
LABORATORIES


Carton Copy  
0.5 gm.

Dispensing Directions: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

STORE IN COOL PLACE

Lot No.

Label Copy  
25 gm.



25 gm.

### MITOX OINTMENT

An aid in the treatment of bacterial infections of the external ear canal associated with ear mite infestation

#### Active Ingredients

Neomycin base (as the sulfate) .....	0.5%
Sulfacetamide .....	9.0%
Carbaryl (1-Naphthyl N-methylcarbamate)* .....	1.0%
Tetracaine hydrochloride .....	0.5%
Mineral oil .....	59.1%
Inert Ingredients .....	29.9%
	100.0%

\*Sevin

**Directions:** If severe accumulations of ceruminous material are present, clean ear before treatment with a ceruminolytic agent. Follow this cleaning with treatment with Mitox, as directed. Fill lower section with Mitox and massage gently to distribute evenly. For infections, repeat as needed. For ear mites, continue treatments at 10 day intervals until controlled.

**NOTE:** After using, retract plunger 1/16 inch to stop the flow of ointment before replacing cap.

Store in cool place

For Veterinary Use Only


Lot No. .

EPA Est. 622-NB-1

EPA Reg. No. 622-38

**CAUTION:** Keep out of reach of children. Harmful if swallowed.

Norden Laboratories, Inc., Lincoln, Nebr. 68501, U.S.A.

 NORDEN  
LABORATORIES

Carton Case  
6-25 cc. syringes

Front and back Panels


6-25 cc. syringes

#### MITOX OINTMENT

An aid in the treatment of bacterial infections of the  
external ear canal associated with ear mite infestation

CAUTION: Keep out of reach of children. Harmful if swallowed.

For Veterinary Use Only

 NORDEN  
LABORATORIES



Carton Copy  
6 - 25 gm. syringes

Right Panel:

6-25 gm. syringes  
Dispense-Pak

#### MITOX OINTMENT

**Directions:** If severe accumulations of ceruminous material are present, clean ear before treatment with a ceruminolytic agent. Follow this cleaning with treatment with Mitox, as directed. Fill lower section with Mitox and massage gently to distribute evenly. For infections, repeat as needed. For ear mites, continue treatments at 10 day intervals until controlled.

**NOTE:** After using, retract plunger 1/16 inch to stop the flow of ointment before replacing cap.

EPA Reg. No. 622-38

EPA Est. 622-NB-1

Norden Laboratories, Lincoln, Nebr. 68501, U.S.A.

Carton Copy  
6-25 gm. syringes

Left Panel:

6-25 gm. syringes  
Dispense-Pak

MITOX OINTMENT

Active Ingredients

Neomycin base (as the sulfate) .....	0.5%
Sulfacetamide .....	9.0%
Carbaryl (1-Naphthyl N-methylcarbamate)* .....	1.0%
Tetracaine hydrochloride .....	0.5%
Mineral Oil .....	59.1%
Inert Ingredients .....	29.9%
	<u>100.0%</u>

\*Sevin

Lot No.