

USE DIRECTIONS:

Use at 2 ounces per gallon. This product will be effective against *Pseudomonas aeruginosa* PRD-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use DETER-SAN-I with a cloth or mop and apply to walls, floors and other hard surfaces.

If used for sanitization of food equipment or food contact items, limit the level of active quaternary to 200 ppm. (2.8 ounces per 5 gallons) NO RINSE REQUIRED.

For heavy soil or organic matter, a precleaning step is recommended.

ACCEPTED

2-8-73

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 623-32

DETER-SAN-I

CLEANER DEODORIZER **DISINFECTANT FUNGICIDE**
 Designed for Hospital and Institutional Use

Phenol Coefficients:

Guaranteed	Minimum
<i>S. aureus</i>	42
<i>S. typhosa</i>	27

ACTIVE INGREDIENTS:

n-Alkyl (C ₁₄ 60%, C ₁₆ 30%, C ₁₂ 5%, C ₁₈ 5%)	
dimethyl benzyl ammonium chlorides	2.25%
n-Alkyl (C ₁₂ 68%, C ₁₄ 32%) dimethyl ethylbenzyl ammonium chlorides	2.25%
Sodium Carbonate	3.00%
INERT INGREDIENTS	92.50%
	100.00%

Rinse empty container thoroughly with water and discard.

NET VOLUME: 1 gallon. EPA REG. No. 623-32

DANGER

Keep Out of Reach of Children.
 See right side panel for additional precautionary statements.

UNITED CHEMICAL COMPANY, INC.
 4th & Delaware Kansas City, Mo. 64105

DANGER:

Keep Out of eye damage, eyes, on skin, face shield, Harmful or t, tion of food

FIRST AID:

In case of contact with plenty eyes, call a taminated c

If swallowed milk, egg w not availab Avoid alcoh

NOTE TO P Probable m the use of circulatory s vulsion may

W-I
STANT
DE
al Use

s)
2.25%
2.25%
3.00%
92.50%
100.00%
ater and
. 623-32
en.
onal
Y, INC.
o. 64105

DANGER:

Keep Out of Reach of Children. Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink **promptly** a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DETER-SAN-I is a cleaner, disinfectant and sanitizer designed for institutional use. Such as HOSPITALS, NURSING HOMES, and SCHOOLS where housekeeping will be of prime importance in reducing the hazard of cross-infection from environmental surfaces.

DETER-SAN-I can provide the assurance you need in this area, when used as directed. For use on floors, walls, sink tops, garbage pails, telephones, rest rooms, bed pans and operating rooms. Effective against Staphylococcus aureus, Salmonella choleraesuis and Pseudomonas aeruginosa. Also effective against pathogenic fungi, Trichophyton interdigitale.

"This product when used on environmental, inanimate, hard surfaces at 2 ounces per gallon of water, is effective against Influenza A₂, Herpes Simplex, Adenovirus type 2, and Vaccinia viruses."