


ACCEPTED
 OCT 21 1975
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 62334

**DEDICATED TO DEPENDABILITY
 SINCE 1913**



Q-NARY-10

DISINFECTANT • DEODORIZER • SANITIZER

PHENOL COEFFICIENTS:

S. Aureus.....	92
S. Typhosa.....	60

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	5
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	5

INERT INGREDIENTS:	90
	100%

Q-NARY-10 is a concentrate and should be diluted before using.

NET CONTENTS: _____ EPA Reg. No. 623-34
 EPA EST. 623-MO-01

DANGER
KEEP OUT OF REACH OF CHILDREN
 (See other precautionary and first aid
 statements on left side panel)

Manu
UNITED C

USE DIRE

Household

For sanitizin
 inanimate h
 Locker Room
 ridors, Class
 Apply solutic
 ounce to 4
 ounces to 5
 cus aureus
 heavily soil
 cleaning ste

Hospitals

For floors, wa
 sing Homes,
 Garbage Pail
 at 3 1/2 ounce
 cloth, mop,
 will be e
 aerugiosa f
 taminated
 recommende

Food Proces

For sanitiza

Cold Disinfection

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1 1/2 ounce per gallon of water for 10 minutes.

FOR ROOM FOGGING

DIRECTIONS:

To minimize the danger of cross-contamination from environmental surfaces apply solution as a fog described below, before applying standard routine terminal cleaning and disinfecting practices. 1. Remove all human, animal, and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount BURGESS MODEL NO. 1601 MISTER or other fogging device delivering equivalent spray at 3 RPM. 5. Fill sprayer reservoir with 1.5 oz. to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed

surfaces. 8. Wait 2 hours before entering room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN:

Probably mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard

ED
 1 1975
 RAL INSECTICIDE
 ODENTICIDE ACT
 OISON REGISTER-
 2334



Q-NARY-10

DISINFECTANT • DEODORIZER • SANITIZER

PHENOL COEFFICIENTS:

S. Aureus.....	92
S. Typhosa.....	60

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	5%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	5%

INERT INGREDIENTS:	90%
	100%

Q-NARY-10 is a concentrate and should be diluted before using.

NET CONTENTS: _____ EPA Reg. No. 623-34
 EPA EST. 623-MO-01

DANGER

KEEP OUT OF REACH OF CHILDREN

(See other precautionary and first aid statements on left side panel.)

Manufactured and Distributed by:
UNITED CHEMICAL COMPANY, INC.

USE DIRECTIONS:

Household

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce to 4 gallons of water. Disinfect with 3 ounces to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use at 3½ ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level Q-Nary-10 will be effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Food Processing Equipment

For sanitization of previously cleaned food

OFFICE AND PLANT: 5050 East 52nd Street
 Kansas City, Mo. 64130
 (816) 921-5050

processing equipment and food utensils dilute 1 ounce per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, No Potable Water Rinse is required.

Dairies

To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce to 4 gallons of water (200 ppm). At this level, No Potable Water Rinse is required. Follow recommendations of local Health Board.

Restaurant and Bar Rinse

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce to 4 gallons of water. **No Terminal Potable Water Rinse Required.**

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A², Herpes Simplex, Adenovirus type 5, and Vaccinia Viruses.