

CEPTED

1975

FEDERAL INSECTICIDE
AND ROENTICIDE ACT
EPA REG. NO. 623-38



ring room
equipment
g use.

Corrosive.
Do not get
t eyes and
harmful if
ood.

h eyes or
5 minutes.
d wash all
f swallow-
ution, or if
antities of

raindicate
gainst cir-
n and con-

with water

UCO MINT DISINFECTANT

DISINFECTANT • DEODORIZER • SANITIZER

PHENOL COEFFICIENTS:

S. Aureus	92
S. Typhosa	60

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	5%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	5%
Isopropyl Alcohol.....	2%
Methyl Salicylate.....	1%

INERT INGREDIENTS:	87%
	100%

Uco Phene Disinfectant is a concentrate and should be diluted before using.

NET CONTENTS: _____ EPA Reg. No. 623-38
EPA EST. 623-MO-01

DANGER

KEEP OUT OF REACH OF CHILDREN

(See other precautionary and first aid statements on left side panel.)

Manufactured and Distributed by:

UNITED CHEMICAL COMPANY, INC.

OFFICE AND PLANT: 5050 East 52nd Street
Kansas City, Mo. 64130
(816) 921-5050

USE DIRECTIONS:

Household

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce to 4 gallons of water. Disinfect with 3 ounces to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use at 3½ ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level UCO Phene Disinfectant will be effective against Pseudomonas aeruainosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A², Herpes Simplex, Adenovirus type 2, and Vaccinia Viruses.

IF THIS PRODUCT
IS USED TO TREAT
FOOD PROCESSING
EQUIPMENT OR
FOOD UTENSILS,
FOOD CONTACT SURFACES
MUST BE SCRUBBED WITH
DETERGENT AND RINSED
WITH POTABLE WATER
PRIOR TO RE-USE.