



ETA-TROL

Disinfectant and Deodorant

ACTIVE INGREDIENTS: Beta Terpineol 40%, Vegetable Oil Soap 10%,
Isopropyl Alcohol 40%
INERT INGREDIENT: Water 50%

FOR MATERIAL CONTROL

A powerful, pleasant disinfectant, deodorant, and germicide. Guaranteed to be 3 times stronger than pure Carbolie Acid when tested against *S. typhosa* (typhoid germ) at 20°C in the A.O.A.C. Phenol Coefficient Method. Because of its pleasant lilac odor ETA-TROL is excellent for use in Hotels, Motels, Schools, Institutions, Office Buildings, Hospitals, Sanatoriums, as a disinfectant and deodorant.

ETA-TROL

Instantly soluble in water in which it forms a creamy milk-white emulsion which does not separate on standing. When mixed with water in recommended dilutions, does not stain the walls, floors, or clothes and leaves a pleasant and agreeable odor of Lilac.

FOR INTERNAL USE ONLY

For use in the treatment of typhoid fever, typhoid pneumonia, typhoid meningitis, typhoid enteritis, typhoid colitis, typhoid hepatitis, typhoid nephritis, typhoid dermatitis, typhoid neuritis, typhoid myelitis, typhoid osteomyelitis, typhoid arthritis, typhoid sinusitis, typhoid otitis media, typhoid otitis externa, typhoid conjunctivitis, typhoid keratitis, typhoid iridocyclitis, typhoid glaucoma, typhoid cataracts, typhoid strabismus, typhoid amblyopia, typhoid blindness, typhoid deafness, typhoid dumbness, typhoid paralysis, typhoid convulsions, typhoid coma, typhoid death.

DOSE

For the treatment of typhoid fever, 4 ounces of ETA-TROL to each gallon

of water. For the treatment of typhoid pneumonia, typhoid meningitis, typhoid enteritis, typhoid colitis, typhoid hepatitis, typhoid nephritis, typhoid dermatitis, typhoid neuritis, typhoid myelitis, typhoid osteomyelitis, typhoid arthritis, typhoid sinusitis, typhoid otitis media, typhoid otitis externa, typhoid conjunctivitis, typhoid keratitis, typhoid iridocyclitis, typhoid glaucoma, typhoid cataracts, typhoid strabismus, typhoid amblyopia, typhoid blindness, typhoid deafness, typhoid dumbness, typhoid paralysis, typhoid convulsions, typhoid coma, typhoid death.

For the treatment of typhoid fever, 4 ounces of ETA-TROL to each gallon of water. For the treatment of typhoid pneumonia, typhoid meningitis, typhoid enteritis, typhoid colitis, typhoid hepatitis, typhoid nephritis, typhoid dermatitis, typhoid neuritis, typhoid myelitis, typhoid osteomyelitis, typhoid arthritis, typhoid sinusitis, typhoid otitis media, typhoid otitis externa, typhoid conjunctivitis, typhoid keratitis, typhoid iridocyclitis, typhoid glaucoma, typhoid cataracts, typhoid strabismus, typhoid amblyopia, typhoid blindness, typhoid deafness, typhoid dumbness, typhoid paralysis, typhoid convulsions, typhoid coma, typhoid death.

For the treatment of typhoid fever, 4 ounces of ETA-TROL to each gallon of water. For the treatment of typhoid pneumonia, typhoid meningitis, typhoid enteritis, typhoid colitis, typhoid hepatitis, typhoid nephritis, typhoid dermatitis, typhoid neuritis, typhoid myelitis, typhoid osteomyelitis, typhoid arthritis, typhoid sinusitis, typhoid otitis media, typhoid otitis externa, typhoid conjunctivitis, typhoid keratitis, typhoid iridocyclitis, typhoid glaucoma, typhoid cataracts, typhoid strabismus, typhoid amblyopia, typhoid blindness, typhoid deafness, typhoid dumbness, typhoid paralysis, typhoid convulsions, typhoid coma, typhoid death.

In a TYPHOID SICK-ROOM, in the disinfecting of beds, clothes, utensils, room surfaces; the sanitary removal of typhoid stools and the sanitizing of hands of those attending typhoid patients, ETA-TROL should be used in the proportion of four ounces in each gallon of water.

For the treatment of typhoid fever, 4 ounces of ETA-TROL to each gallon of water. For the treatment of typhoid pneumonia, typhoid meningitis, typhoid enteritis, typhoid colitis, typhoid hepatitis, typhoid nephritis, typhoid dermatitis, typhoid neuritis, typhoid myelitis, typhoid osteomyelitis, typhoid arthritis, typhoid sinusitis, typhoid otitis media, typhoid otitis externa, typhoid conjunctivitis, typhoid keratitis, typhoid iridocyclitis, typhoid glaucoma, typhoid cataracts, typhoid strabismus, typhoid amblyopia, typhoid blindness, typhoid deafness, typhoid dumbness, typhoid paralysis, typhoid convulsions, typhoid coma, typhoid death.

HARMFUL IF SWALLOWED

