


E-Z FLO 80% SUPERFINE SULFUR

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS	
Sulfur	80.0%
INERT INGREDIENTS	20.0%
	<hr/>
	100.0%



DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

FOLLOW DIRECTIONS CAREFULLY

Application: Apply at rate of 1/2 to 2 pounds per tree, depending on size of tree. Dust thoroughly to obtain good coverage on fruit and foliage.

APPLES: For control of apple scab and powdery mildew. Supplement early scab sprays at petal fall and during early covers. Also may be used to combat late scab infection during weather periods favorable for scab development. **CAUTION:** Do not use Sulfur within 10 to 14 days of an oil spray.

PEACHES: For control of peach scab and brown rot (Blossom blight). Apply during bloom, and first cover (about three weeks after petal fall). Repeat about one month before harvest and continue at weekly intervals.

PLUMS: For control of brown rot (blossom blight) and leaf spot. Apply during bloom period, petal fall, shuck split, first, third and fourth cover sprays.

SWEET CHERRIES: For control of brown rot and leaf spot. Use in a complete dust schedule.

CAUTION

Avoid application during or just prior to periods of extremely high temperatures and humidity.