


E-Z-FLO COPPER 7 DUST



CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Copper (In Tribasic Copper Sulphate)
expressed as Metallic Copper 7.00%

INERT INGREDIENTS: 93.00%

Total 100.00%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

HARMFUL IF SWALLOWED

This product is toxic to fish. Keep out of any body of water.

Do not apply when weather conditions favor drift of dust from crop areas treated.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED.

POTATOES — For control of early and late blight. Apply 30-35 pounds per acre beginning when plants are 2-5 inches tall. 3 to 5 late applications are usually necessary. (Weather conditions will determine frequency of applications)

CELERY -- For control of early and late blight. Apply at rate of 30-35 pounds per acre starting when plants are small. Continue at 5-7 day intervals.

SUGAR BEETS — For control of Cercospora leaf spot. Apply at 25-35 pounds per acre, starting before disease threatens and continue at 10 day intervals as often as necessary for control.

TOMATOES — For control of late blight. Apply 25 to 35 pounds per acre at 7 day intervals beginning when plants are 2 to 5 inches tall. Apply as often as necessary.