

E-Z-Flo Methoxychlor 5 Dust

CAUTION:

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

*Methoxychlor Technical.....5.00%
*Equivalent to 4.4% 2,2-bis (p-methoxyphenyl)-1,1,1-Trichloroethane
and 0.6% other isomers and related compounds.

INERT INGREDIENTS:.....95.00%
Total 100.00%

CAUTION

Avoid inhalation and contact with skin. Wash with soap and water after using. Keep away from children, pets, livestock and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.


This product is toxic to bees and should not be applied when bees are actively visiting the area.

Destroy Container When Empty. Never Re-use.

51-30

EPA Est. 635-M1-1; OH-

E.P.A. Reg. No. 635-16



DIRECTIONS FOR USE

Apply when insects first appear and follow with repeated applications at intervals of 7 to 10 days until insects disappear. Apply at the rate of 20 to 35 pounds per acre unless otherwise indicated. Apply for thorough coverage.

Remove residues on edible portions of vegetables at harvest by washing, brushing or other effective means.

DTH - Indicates Days To Harvest limitations. Do not apply within the indicated DTH.

ASPARAGUS- Asparagus beetles: 3 DTH unless washed or blanched.

BEANS- Mexican bean beetle, leafhoppers, Japanese beetle, Bean leaf beetle: 3 DTH.

BEETS (table)- Blister beetles and flea beetles: Apply 25 to 30 pounds per acre and repeat at 7-10 day intervals as necessary. Do not use on sugar beets. 7 DTH, 14 DTH if tops are used for food or feed.

BLACK-EYED PEAS- Cowpea curculios: Apply 30 to 35 pounds per acre. Do not feed treated vines to livestock. 3 DTH.

BROCCOLI- Cabbage worm and flea beetles: 14 DTH.

BRUSSEL SPROUTS- Cabbage worm and flea beetles: 14 DTH.

CABBAGE- Imported cabbage worm, flea beetles: 3 DTH.

CARROT- Leafhoppers. Apply 35 to 40 pounds per acre: 7 DTH, 14 DTH if tops are used for food or feed.

CAULIFLOWER- Imported cabbage worm and flea beetle: 7 DTH.

CORN- Fall armyworm, flea beetles, Japanese beetle, and rose chafers: 7 DTH.

CUCUMBER, MELONS, SQUASH- Cucumber beetles, leafhoppers and Melon worms: 1 DTH.

EGGPLANT- Flea beetles and leafhoppers: 1 DTH.

LETTUCE- Leafhoppers: 14 DTH.

PEANUTS- Velvetbean caterpillar: 7 DTH.

PEAS- Pea weevil: 7 DTH.

PEPPERS- Flea beetles and rose chafers: 1 DTH.

POTATOES- Colorado potato beetle, Leafhopper, Flea beetle: 0 DTH.

RADISH- Flea beetles: 7 DTH.

RUTABAGA- Flea beetles: 7 DTH.

SOYBEANS- Mexican bean beetles, Soybean caterpillar, Velvetbean caterpillar: 7 DTH.

SPINACH- Flea beetles, Leafhoppers, Diabrotica beetles: 14 DTH.

TOMATOES- Blister beetles, fall armyworm, flea beetles, Japanese beetle, tomato fruit worms: 1 DTH.

TURNIP- Flea beetles: 7 DTH, 14 DTH if tops are used for food or feed.

PUMPKIN - Cucumber beetle and squash vine bore: 1 DTH.