

[635-41]

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Rotenone **4.0%**

Other Cube Resins **8.0%**

INERT INGREDIENTS: **88.0%**

ACCEPTED

635-41

JUL 10 1974

UNLESS OTHERWISE SPECIFIED
FOR MORE INFORMATION CONTACT
EPA REGIONAL OFFICE
EPA OFFICE OF PESTICIDE REGISTRATION
10 WILLIAMSON GARDEN

100.0%

*Net Contents per Manufacturer
On Container*

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin. Avoid inhalation of dust or spray mist. Wash thoroughly after using.

This product will kill fish. Keep out of lakes, ponds, and streams.

Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas.

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED: Do not apply within one day of harvest. Sift material into spray tank when it is about half full of water. Use 75 to 125 gallons water per acre depending on size of plants. Apply thoroughly when insects first appear to both upper and under leaf surfaces. Follow with repeated applications at intervals of about seven days and/or after rains until insects disappear.

ASPARAGUS: For control of asparagus beetle apply 4-6 lbs. per acre. Apply as often as necessary for control to spears or fern growth.

BEANS: For control of Mexican bean beetle apply 4-6 lbs. per acre. Apply with high pressure spray equipment. Make sure material is covering underside of bean leaves. Repeat as necessary.

CABBAGE, CAULIFLOWER: For control of cabbage looper and Imported cabbageworm apply 6.5 to 8.0 lbs. per acre. Start applications early in growing season and continue as necessary, usually at 7-10 day intervals.

PEAS: For control of aphids (except woolly aphids) and pea weevil apply 6.5 to 8.0 lbs. per acre. High pressure equipment is necessary to drive spray through mat of vines. For control of pea weevil repeat at 4-5 day intervals until harvest.

CUCUMBER, MELON, SQUASH: For control of striped and spotted cucumber beetles apply 7.5 to 10.0 lbs. per acre. Start applications when plants emerge from ground and continue at weekly intervals.