

E-Z FLO DITHIANE 6 DUST

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

*Zinob (zinc ethylene bisdithiocarbamate)
(Zinc Metallic Equivalent 96%) 3.9%

INERT INGREDIENTS: 96.1%

100.0%

*Equivalent to 6% Dithane Z-78

ACCEPTED

MAY 28 1971

IN ACCORDANCE WITH THE FEDERAL INSECTICIDE
ACT AND THE FEDERAL FERTILIZER ACT
FOR THE YEAR 1971, THIS PRODUCT IS REGISTERED
UNDER E.P.A. REG. NO. 635-44. SUBJECT
TO ATTACHED COMMENTS.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

Use Only as Directed: Begin application at first sign of disease and repeat at 7 to 10 day intervals to keep new growth thoroughly covered. Apply 20 to 25 pounds per acre.

ONIONS: Botrytis Leaf Blight, Downy Mildew, Purple Blotch. Do not apply within 7 days of harvest on green onions.

TOMATOES: Anthracoze, Early Blight, Late Blight, Cladosporium Gray Mold, and Stemphylium Leaf Spot.

CUCUMBERS, MELONS and SQUASH: Alternaria (Macrosporium) Blight, Anthracnose, Downy Mildew, and Gummy Stem Blight. Direct dust to the underside of the leaves.

CAUTION

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN.
MAY BE HARMFUL IF INHALED OR SWALLOWED.

Avoid breathing dust. Avoid contact with skin, eyes and clothing. In case of contact flush with plenty of water; for eyes, get medical attention. Keep containers closed. Store in dry place. Keep away from fire and sparks.