

ACCEPTED
MAY 14 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. ()

E-Z FLO
COPPER, METHOXYCHLOR 5-3 DUST

CAUTION:
 KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Copper (In Basic Copper Sulfate) expressed
 as metallic.....5.0%

*Methoxychlor, Technical..... 3.0%

INERT INGREDIENTS:.....92.0%
 100.0%

*Equivalent to 2.64% 2,2-bis (p-methoxyphenyl)-1,1,1-
 Trichloroethane and .36% other isomers and related compounds.

CAUTION

Avoid breathing dust. Avoid contamination of feed and food stuffs.
 Avoid skin contact. In case of skin contact, wash with soap and water.

Continued on side/back panel-



54-22 EPA Est 635-MI-1; OH-1 E.P.A.Reg. No. 635-103

DIRECTIONS FOR USE

CUCUMBERS, MELONS, PUMPKINS and SQUASH: For control of striped and spotted cucumber beetles, bacterial wilt (indirectly by control of cucumber beetles) angular leaf spot, downy mildew, scab, and *Macrosporium* leaf spot. Apply 35 to 40 pounds per acre depending on plant size and disease and insect conditions. Make first application as soon as the young plants appear above ground and follow with repeated applications at intervals of one week and/or after rains until insects disappear and danger of disease infection is past. Do not apply within 1 day of harvest.

CAUTION

Continued from front panel-

Highly toxic to fish. Do not use where any body of water will be contaminated by direct application or runoff. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Destroy Container When Empty. Never Re-use.