

# E-Z FLO VEGA-GUARD

**CAUTION:**  
KEEP OUT OF REACH OF CHILDREN

**ACTIVE INGREDIENTS:**

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate).....5.00%  
Zineb (zinc ethylene bisdithiocarbamate).....10.00%

**INERT INGREDIENTS:** .....85.00%  
100.00%

**CAUTION**

Harmful if swallowed. May cause irritation of nose and throat. Do not breathe dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

See other Precautions on the side/back panel-

55-23


EPA Est. 635-M1-1; OH-1

E.P.A. Reg.No. 635-112

**DIRECTIONS FOR USE**

Apply 20 to 30 pounds as a dust per acre unless indicated otherwise. Begin when diseases threaten or insects first appear using the lower rate when plants are small and increasing the rate to 30 lbs. per acre as plants reach maturity. Repeat at 7 to 10 day intervals, more frequently under severe disease or insect conditions.

BEANS (lima, snap)- For the control of Mexican Bean Beetle, Leafhoppers, Japanese Beetle, Anthracnose, Downy Mildew and Rust: Apply 25-30 pounds per acre. Do not apply within 7 days of harvest.



**DIRECTIONS FOR USE** (Continued)

BEETS- For the control of Downy Mildew and Leaf Spots: Do not apply within 7 days of harvest.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, and CAULIFLOWER- For the control of Aphids, Imported Cabbage Worm, Cabbage Loopers, Downy Mildew and Leaf Spots: Apply 25-30 pounds per acre. Do not apply within 7 days of harvest.

CARROTS- For the control of Aphids, Leafhoppers, Early and Late Blights: Do not apply within 7 days of harvest.

CUCUMBER, MELONS (cantalope, cassabas, crenshaws, honeydews, honeyballs, muskmelon, persian melons, watermelons) and SQUASH- For the control of Aphids, Spider Mites, Pickleworms, Alternaria Blight, Anthracnose, Blossom Blight, Downy Mildew and Gummy Stem Blight: Begin when plants emerge or at first sign of disease or insects. Apply 25-30 pounds per acre. Do not apply within 5 days of harvest.

EGGPLANT- For the control of Alternaria leaf spot (early blight) begins in plant bed. Anthracnose, Cercospora leaf spot, downy mildew, Phomopsis and other fruit rots. Phytophthora (late blight). Do not apply within 3 days of harvest.

PEPPERS- For the control of Anthracnose, Downy Mildew, Cercospora and other leaf spots, Phytophthora and other fruit rots: Do not apply within 3 days of harvest.

TOMATOES- For the control of Aphids, Spider Mites, Russet Mites, Drosophila, Anthracnose, Early and Late Blight, Gray Mold (Cladosporium), Gray Leaf Spot (Stemphylium), Nail Head Spot, Septoria Leaf Spot, and Southern Blight (Sclerotium): Begin when first fruit clusters are well formed or when disease or insects first appear. Apply 25-30 pounds per acre. Do not apply within 5 days of harvest.

**CAUTION**

Continued from panel on left-

Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Destroy Container When Empty. Never Re-use.

Net Content: 1.5 lbs. Manufacturer: ...