

E-Z FLO CHLORDANE 40 WP

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Technical Chlordane* 40.0%

INERT INGREDIENTS: 60.0%

100.00%

*Equivalent to 24% octachloro-4, 7-methanotetrahydroindane and 16% related compounds.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

Measure the specified amount of this product then wash through filler screen in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants. Emulsifiable concentrates should be thoroughly emulsified with at least half of total water before adding this product.

RECOMMENDATIONS FOLIAGE TREATMENT

Unless otherwise specified, use 2 1/2 to 3 lb. Chlordane 40 WP per acre in sufficient water for thorough coverage.

BEANS: Armyworms (fall, Southern, yellow-striped), cucumber beetles (12-spotted, banded), flea beetle, green stink bug. Do not apply to green or snap beans after pods start to form.

CARROTS: Carrot rust fly. Do not use tops for food or feed.

CELERY: Carrot rust fly, cutworms. Apply preplant or at planting time only.

COLE CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower) - Cabbage worm, flea beetle, green stink bug, harlequin bug. Do not apply after heads or edible parts start to form.

CORN (SWEET): Silk fly. Use 2 1/2 lb. per acre making first application before silking. Make 1 to 3 applications at 5-day intervals if needed. **CORN:** Corn rootworm, white-fringed beetle larvae. Use 5 to 10 lb. per acre in sufficient water for thorough coverage of infested area or areas to be protected. Apply as a preplant or at planting only. Do not feed treated forage to dairy animals or animals being finished for slaughter.

CORN (FIELD AND SWEET): Broadcast application: Make broadcast application of 6.25-10 pounds per acre for control of the soil insect complex, including wireworms, white grubs, seed corn beetle, seed corn maggot and larvae of green June bug, Asiatic garden beetle, Japanese beetles and white fringed beetles. For the control of corn rootworms (Western, Southern, Northern), cutworms, cornfield ants, and corn root aphids apply 5-10 pounds per acre.

Apply after the field is plowed. Disk in promptly. Use the higher rate on heavier soils and the lower rate on light sandy soils. Use the higher rates for heavy infestation of cutworms.

Row application: To control corn rootworms (Western, Northern, Southern), cornfield ants and corn root aphids apply 2.5 pounds per acre at planting time in the furrow, behind the planter shoe and ahead of the press wheel. To control wireworms and white grubs, apply 6.25 pounds per acre in the row at planting time.

Consult your State Agriculture Extension Service for detailed recommendations.

EGGPLANT: Flea beetle, serpentine leafminer. Do not apply after edible parts start to form.

ONIONS: Thrip. Do not apply to green or spring onions.

PEAS: Flea leafminer. Do not apply within 15 days of harvest. Do not feed treated vines to dairy animals or animals being finished for slaughter.

STRAWBERRIES: Crown borer, field crickets, lesser cornstalk borer, spittle-bug. Apply before blossoms appear. Do not apply after fruit begins to form.

TOMATOES: Field crickets, leaf-footed plant bug, leaf miner, stink bug. Do not apply after fruit begins to form.

SMALL GRAINS (Barley, Oats, Rye, Wheat): Ants, crickets, grasshoppers. Do not apply after the heads start to form. Do not allow grazing in treated fields.

SOIL INSECT CONTROL

WHITE GRUBS: Potatoes, strawberries and sugar beets. Apply 10 lb. maximum in 100 gal. water per acre as a pre-plant treatment to soil surface. Disk into top 3 or 4 inches of soil.

WIREWORMS: Beans, beets, broccoli, cabbage, cauliflower, corn, onions, potatoes, strawberries, sugar beets, sweet potatoes and turnips. Make pre-planting treatment to soil surface applying 10 lb. maximum in 100 gal. water per acre. Disk into top 4 to 6 inches of soil.

CUTWORMS: Beans, corn, eggplant, lettuce, onions, pepper, potatoes, and tomatoes. Apply 2 1/2 lbs. per 100 gal. water (150 gal. per acre) before planting. Apply evenly over soil surface and do not disturb soil for 3-5 days.

MOLE CRICKETS: Broccoli, cabbage, cauliflower, lettuce, strawberries and tomatoes. Apply 2 1/2 to 5 lb. per acre evenly over soil surface before seeding or transplanting, using sufficient water to insure complete coverage.

TERMITES: Apply 10-20 lbs. in 100 gal. water. Apply at rate of 1 gallon per linear foot to a trench dug next to the foundation 1 to 2 feet deep and 8 inches wide. For buildings with deep footings, dig trench 30 in. deep. Do not dig trench below the level of the footing. Inspect area yearly to determine possible reinfestation.

NOTE: When necessary to make foliage sprays to control ants on the crops listed on this label observe restrictions and do not apply on forage crops which are to be fed to dairy animals or animals being finished for slaughter.

ACCEPTED
Aug 26, 1971
UNDER THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
OF 1947 AS AMENDED
EPA ECONOMIC POISON REGISTER
E.P. NO. 635-120

Net Contents and Manufacturer
On Container

CAUTION

Keep out of reach of children. If Swallowed, Avoid Breathing Spray Mist. Contact With Skin Can Cause Toxic Symptoms. In Case of Contact With Skin, Wash With Soap and Water. Avoid Contamination of Feed and Foodstuffs.

OBSERVE THE FOLLOWING USE PRECAUTIONS:

Observe use limitations given on the label for specific crops.

Do not apply on cucurbits (cantaloupe, cucumber, squash, melon) as injury to plants may result. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply under conditions favoring drift to adjoining food, fibre or pasture crops, or to areas occupied by unprotected humans, domestic or wild animals. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Highly toxic to fish and wildlife. Do not apply under conditions where contamination of any body of water can occur directly, by drift or runoff. Birds feeding on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

Consult your State Agricultural Experiment Station for additional information consistent with label directions and cautions, as the timing, number and rate of applications needed will vary with local conditions.

Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).