

E-Z-FLO MANEB 7 DUST

CAUTION:
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT:

Maneb (Manganese Ethylenebisdithiocarbamate)	7.0%
INERT INGREDIENTS:.....	.93.0%
Total	100.0%

CAUTION

May cause irritation of eyes, nose, throat, and skin. May be harmful if inhaled or swallowed. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water; for eyes, get medical attention.

This product is toxic to fish. Keep out of any body of water.

Destroy container when empty. Never re-use.

IMPORTANT: Never allow dust to become wet during storage as it will reduce the effectiveness of the dust and it will create flammable vapors. Keep container closed when not in use.

52-27

EPA Est 635-MI-1; OH-1

EPA Reg. No. 635-129

Net Contents and Manufacturer
On Container

DIRECTIONS FOR USE

BEANS (snap or string) - Anthracoze, Downy Mildew, Rust: Apply 15 to 45 pounds/acre. Begin when diseases threaten. Repeat at 5 to 7 day intervals. No time limitation on dry beans; do not apply within 4 days of harvest on succulent beans.

BROCCOLI- Alternaria leaf spot, Downy mildew: Apply 17.2 to 57 pounds/acre. Begin when diseases threaten. For downy mildew begin in plant bed and repeat at 3 day intervals. In field, repeat at 7 to 14 day intervals. If applied within 3 days of harvest, remove residues by washing or trimming.

BRUSSELS SPROUTS, CAULIFLOWER, AND KOHLRABI- Alternaria leaf spot, Downy mildew: Apply 17 to 37 pounds/acre. Begin when diseases threaten. Repeat at 3 day intervals in plant beds and at 7 to 10 day intervals in field.

CABBAGE- Alternaria leaf spot, Downy mildew: Apply 15 to 37 pounds/acre. Begin 7 to 10 days after planting. Repeat at 3 to 5 day intervals in plant bed and field. Do not apply within 7 days of harvest.

CELERY- Early blight (Cercospora), Late blight (Septoria): Apply 25 to 40 pounds/acre. Begin when plants emerge in plant beds. Repeat at 3 to 5 day intervals in plant beds and at 7 to 10 day intervals after plants are set in field. Do not apply within 14 days of harvest. Remove excess residues by stripping, trimming, and washing.

CUCUMBERS- Alternaria (Macrosporium) leaf spot, Angular leaf spot, Anthracnose, Downy mildew, Pythium fruit rot: Apply 17 to 40 pounds/acre. Begin when diseases threaten or plants begin to run. Repeat at 7 to 10 day intervals or at 3 to 5 day intervals under severe disease conditions. Do not apply within 5 days of harvest.

EGGPLANT- Anthracoze, Early blight (Alternaria), Phomopsis blight or Fruit rot: Apply 15 to 40 pounds/acre. Begin when fruit starts to form. Repeat at 7 to 10 day intervals.

ENDIVE, LETTUCE- Downy mildew: Apply 26 to 45 pounds/acre. Begin when disease appears. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce and endive by washing or other effective means.

MELONS (Cantaloupes, Casabas, Crenshaws, Honeydews, Honey Balls, Muskmelons, Persian Melons and Watermelons)- Anthracoze, Cercospora leaf spot, Downy mildew, Gummy stem blight: Apply 16 to 40 pounds/acre. Begin at late bunching, when vines begin to run or when disease first appears. Repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

ONIONS- Botrytis leaf blight (blaf), Downy mildew, Purple blotch: Apply 25 to 40 pounds/acre. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

PEPPERS- Anthracoze, Cercospora leaf spot (foryeye spot), Ripe rot: Apply 15 to 40 pounds/acre. Begin when disease threatens. Repeat at 7 to 10 day intervals.

POTATOES- Early and Late blights: Apply 18 to 56 pounds/acre. Begin when plants are 2 to 6 inches high. Repeat at 5 to 10 day intervals, more often under severe late blight conditions.

TOMATOES (greenhouse and field)- Cladosporium leaf mold, Early and Late blights, Gray leaf spot (Stemphylium), Septoria leaf spot: Apply 18 to 57 pounds/acre. Begin at first fruit cluster and repeat at 7 to 10 day intervals. Anthracoze: Apply 42 pounds/acre. Begin when first fruit forms. Repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

TOBACCO- Anthracoze: Apply 26 to 45 pounds/acre. Begin after plants are set in field. Repeat as necessary. Blue mold: Apply 0.6 to 1.8 pound/100 square yards of plant bed. Begin when plants are about the size of a dime or when blue mold is first reported in area. Repeat twice weekly until plants are set in field. For disease control under field conditions, use 25 to 45 pounds/acre. Repeat as necessary.

ACCEPTED
APR 17 1970
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 635-129