

E-Z FLO ARSENATE OF LEAD

ACTIVE INGREDIENTS:

Lead arsenate* (PbHAsO4)	98.0%
INERT INGREDIENTS:	2.0%
	<hr/> 100.0%

*Total arsenic expressed as metallic..... 21.14%
 Arsenic in water soluble form, expressed
 as metallic, Not more than 0.35%
 Lead expressed as metallic 58.00%

BURN BAG WHEN EMPTY — BURY ASHES.

DIRECTIONS FOR USE

FOLLOW DIRECTIONS CAREFULLY:

Always use freshly prepared spray, mixing only the amount needed. On fruits, mix an equal amount of hydrated lime or other safener with the required amount of lead arsenate.

APPLES, PEARS, QUINCES: For control of Codling Moth, Plum Curculio, Canker Worm, Red Banded Leaf Roller, Tent Caterpillars and Apple Maggot — use 2 to 3 lbs. per 100 gals. of water. Apply in regular spray schedule from pre-pink through cover sprays as needed for control. Make no application later than 30 days before harvest. Remove excess residue at harvest. Do not graze livestock on treated areas.

PEACH, PLUM, CHERRY: To control Plum Curculio use 2 to 3 lbs. per 100 gals. water. Make 2 to 3 applications at 8 to 10 day intervals, beginning at petal fall or shuck split. For Cherry Fruit Flies use 2 lbs. arsenate of lead plus 3 lbs. hydrated lime per 100 gals. water; apply when flies first emerge, make second application 10 days later. On peaches and plums do not apply within 30 days of harvest. On cherries do not apply within 30 days for fresh market nor within 14 days of harvest for processing cherries. Remove excess residue at harvest. Do not graze livestock on treated areas.

Not Contents and Manufacturer
 On Container

86-21

ORNAMENTAL TURF, GOLF COURSES, EMBELLISHMENT AND OTHER NON-CROP AREAS

Apply to ornamental turf, golf courses, embellishment and other non-crop areas. Use 300 to 500 lbs. of E-Z FLO ARSENATE OF LEAD per acre. Mix with 100 lbs. of sand or soil and apply with fertilizer spreader. Apply any time ground is moist or after irrigation or spring applications are preferred. Immediately after application water in thoroughly. Retreat annually with 2 to 4 lbs. per 1,000 sq. ft. to prevent worm and grub damage.

OLD FAIRWAYS: Grubs, Earthworms. When there is a heavy infestation, use 300 to 500 lbs. of E-Z FLO ARSENATE OF LEAD per acre. Mix with 100 lbs. of sand or soil and apply with fertilizer spreader. Apply any time ground is moist or after irrigation or spring applications are preferred. Immediately after application water in thoroughly. Retreat annually with 2 to 4 lbs. per 1,000 sq. ft. to prevent worm and grub damage.

NEW LAWNS, GREENS, TEES: Grubs and Earthworms. To grub and worms on greens and tees under construction, mix 10 lbs. of E-Z FLO ARSENATE OF LEAD with a quantity of moist soil or sand and apply to 1,000 sq. ft. immediately after application water in thoroughly. The application should be made after all grading is completed but before seeds or stolons are planted.

NOTE: Rates recommended for turf may damage some ornamentals if applied too near the roots. Do not apply to dichondra or St. Augustine lawns.

CAUTION: Do not apply this product to feed or forage areas for animals.

USE PRECAUTIONS

Do not apply during bloom when honey bees or other pollinators are active. Avoid contact with spray or dust drift on flowers, shrubs, or fruits, vegetables, and forage crops. Keep children and animals off treated areas until chemical has been washed into the soil by sprinkling or rain and the grass is dry.

Destroy empty bags by burning — bury ashes. Wash spray equipment thoroughly after use. Dispose of wash water where it will not be a hazard to humans or animals. Do not use this product on feed or forage to be fed to animals.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates and do not apply closer to harvest than specified. Consult local agricultural authorities for specific information, as the timing, number and rate of applications will vary with local conditions. This product is toxic to wildlife. Birds and other wildlife in treated areas may be killed.

WARNING



POISON



ANTICOR

53-263 589

U.S.D.A. Reg. No. 635-143

100 0%

BURY ASHES

OR USE

... ..

... ..

... ..

... ..

tees under construction, mix 10 lbs. of E-Z FLO ARSENATE OF LEAD with a quantity of moist soil or sand and apply to 1,000 sq. ft. Immediately after application water in thoroughly. The application should be made after all grading is completed but before seeds or stolons are planted.

NOTE Rates recommended for turf may damage some ornamentals, if applied near the roots. Do not apply to dichondra or St. Augustine lawns.


CAUTION Do not apply this product to feed or forage areas for animals.

USE PRECAUTIONS

... ..

Destroy empty bags by burning or bury ashes. Wash spray equipment thoroughly after use. Dispose of wash water where it will not be a hazard to humans or animals. Do not use this product in feed or forage to be fed to animals.

The Federal Food and Drug Administration has set official tolerance for certain pesticide residues in order that residues on food and forage crops will not exceed established tolerances, if only at recommended rates and do not apply closer to harvest than specified. Consult local agricultural authorities for specific information, as the timing, number and rate of applications will vary with local conditions. This product is toxic to wildlife. Birds and other wildlife in treated areas may be killed.



63-263-584

U.S.D.A. Reg. No. 635-143