

E-Z FLO FRUIT GUARD

(HOME GARDEN USE)

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Malathion (O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate)	5.0%
Methoxychlor Technical*	15.0%
Captan (N-Trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	10.0%
INERT INGREDIENTS:	<u>70.0%</u>
	100.0%

*15% Methoxychlor technical is equivalent to 13.2% 2,2-Bis (p-Methoxyphenyl)-1,1,1-Trichloroethane and 1.8% other isomers and reaction products.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

For disease control, begin applications before diseases appear and repeat at 7 - 10 day interval. Use a dependable pressure sprayer or hand duster, applying to top and bottom leaf surfaces and all parts of fruit.

APPLES: Apply 8 oz. as a dust or mix 8 oz. in 4 gals. water per tree for the control of scab, black rot, brook spot, sooty blotch, apple maggot, codling moth, curculio, and tent caterpillars. Apply in delayed dormant, pre bloom, petal fall and 3 to 4 cover sprays. Cover sprays should be at 7-10 day intervals following petal fall. Do not apply within 7 days of harvest. May injure foliage or some varieties.

PEACHES, APRICOTS. Apply 8 oz. as a dust, or mix 8 oz. in 4 gals. of water per tree. For control of brown rot, scab, and curculio,

apply in delayed dormant, pre-bloom, petal fall, shucksplit, and 2 cover sprays at 7-10 day intervals and a pre-harvest spray 3 weeks before harvest. Do not apply within 21 days of harvest.

PEARS: Apply 8 oz. as a dust or mix 8 oz. in 4 gals. of water per tree. For the control of scab, fruit spot, pear slug, and codling moth. Spray at 7-10 day intervals following petal fall. Do not apply within 7 days of harvest.

STRAWBERRIES: Apply 2 oz. as a dust or mix 2 oz. in 1 gal. of water per 200 sq. ft. for the control of botrytis rot, and tarnished plant bug. For root weevil control, application should be made before planting and worked into the top 6-8 inches of soil. Do not apply later than 14 days before harvest.

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Do not breathe dust mixtures. Wear a respirator of a type approved by the U. S. Bureau of Mines when handling this product. Wash thoroughly with soap and water after use. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals.

NOTE TO PHYSICIANS: Malathion may cause cholinesterase inhibition. Atropine is antidotal.

Highly toxic to fish. Do not contaminate any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agent or Extension Service.

Net Contents and Manufacturer On Container