

# E-Z-FLO Parathion 2 DUSTED

**WARNING:**  
KEEP OUT OF REACH OF CHILDREN

**DUSTED**  
MAY 15 1975  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2-0%

ACTIVE INGREDIENT

Parathion (O), O-Diethyl O-p-Nitrophenyl Thiophosphate

INERT INGREDIENTS: 98.0%

## WARNING

May be fatal if swallowed, inhaled or absorbed through the skin. Avoid getting it on the skin, in the eyes or on clothing. Do not breathe dust. Wear a respirator which has been tested by the U. S. Bureau of Mines for parathion protection. Wear waterproof or protective clothing, natural rubber gloves and hat. Keep away from all feed and food products. Wash hands, arms and face with soap and water thoroughly after application or before eating or smoking. Wash contaminated clothing with hot water and soap before re-using. Keep all unprotected persons or animals out of the operating areas or vicinity where there may be danger of drift. Vacated

Continued on side back panel-

53-21 EPA Est. 635-MI-1; OH-1 E. P. A. Reg. No. 635-163  
**DIRECTIONS FOR USE**

- Apply when insects appear and repeat at 7-10 days or as necessary for control. Do not apply within the indicated Days To Harvest (DTH).
- BEANS** - Aphids, amylworm, bean leaf beetle, leafhoppers (including potato leafhoppers), leafminers (including serpentine leafminer), Mexican bean beetle, spider mites (including 2-spotted spider mites), and stink bug. Apply 25 lbs A, 7 DTH.
  - BEEFS** - Aphids, blister beetles, flea beetles, leafhoppers, leafminers, spider mites, and webworms. Apply 25 lbs A, 15 DTH for roots. 21 DTH if tops are to be used for food or feed.
  - BROCCOLI and BRUSSEL SPROUTS** - Aphids (including turnip aphid), amylworm, cucumber beetle, diamond back moth, imported cabbageworm, leafhopper, leafminers (including serpentine leafminer), spider mites (including 2-spotted spider mites), and thrips. Apply 25 lbs A, 7 DTH.
  - CABBAGE** and **AUTUMN CROPS** - Aphids (including turnip aphid), amylworm, cabbage looper, cucumber beetle, diamond back moth, imported cabbageworm, leafhopper, leafminers (including serpentine leafminer), spider mites (including 2-spotted spider mites), and thrips. Apply 25 lbs A, 10 DTH on cauliflower, 10 DTH on cabbage.
  - CARROTS** - Aphids, leafhoppers, and stink bug. Apply 25 lbs A, 15 DTH.
  - CELERY** - Aphids, cabbage looper, celery leafminer, Colorado potato beetle, fleabeetles, leafhoppers (including aster leafhoppers), leafminers (including serpentine leafminer), spider mites (including 2-spotted spider mites), tarnished plant bug, and thrips. Apply 25 lbs A, 21 DTH.
  - CUCUMBERS** - Aphids, amylworm, cabbage looper, diamondback moth, imported cabbageworm, thrips and vegetable weevil. Apply 25 lbs A, 10 DTH.
  - CORN** - Aphids, amylworm, corn earworm, European corn borer, grasshoppers, spider mites and sap beetles. Apply 25 lbs A, 10 DTH for earworms, apply to silks as they appear. Apply to husks at 2-2 1/2 days before harvest. Repeat immediately after rain. For European corn borer, apply when insects appear, starting at the first or third instar stage, or after 20 egg masses are counted per 10 stalks and first eggs are in the third or fourth stage. If 75% of the plants show shot-hole feeding. Repeat at 5-10 day intervals as needed. Consult your state agricultural experiment station for proper timing. 12 DTH.
  - CUCUMBERS and MELONS** - Aphids (including turnip aphid), amylworm, cucumber beetles (including spotted and striped cucumber beetles), diamondback moths, fleabeetle, leafhopper, leafminers (including serpentine leafminer), and thrips. Apply 25 lbs A, 10 DTH on cucumbers, 7 DTH on melons.
  - PLANTAIN and PEPPER** - Aphids, Colorado potato beetle, flea beetles, leafminers (including serpentine leafminer), and thrips. Apply 25 lbs A, 10 DTH for pepper maggot control on peppers. Apply 20 lbs A, 15 DTH.
  - POTATOES** - Aphids, amylworm, Colorado potato beetle, flea beetles, grasshoppers, leafhoppers (including 2-spotted spider mites), and thrips. Apply 25 lbs A, 10 DTH for potato beetle, grasshoppers, and thrips. Apply 25 lbs A, 15 DTH for leafhoppers, spider mites, and vegetable weevil.

- PEAS** - Aphids (including wireworms and cutworms), grasshoppers, leaf beetles, leafhoppers, leafminers (including serpentine leafminer), Colorado potato beetle, flea beetles, grasshoppers, leafhoppers, spider mites (including 2-spotted spider mites), and thrips. Apply 25 lbs A, 10 DTH for peas, 15 DTH for pea storage.
- POTATOES** - Aphids, amylworm, Colorado potato beetle, flea beetles, grasshoppers, leafhoppers (including 2-spotted spider mites), and thrips. Apply 25 lbs A, 10 DTH for potato beetle, grasshoppers, and thrips. Apply 25 lbs A, 15 DTH for leafhoppers, spider mites, and vegetable weevil.
- PUMPKINS** - Aphids, cucumber beetles, serpentine leafminer, meadow sparrow worm, squash vine borer, and thrips. Apply 25 lbs A, 10 DTH.
- RAISHES** - Aphids, leafminers. Apply 25 lbs A, 15 DTH.
- SPINACH** - Aphids, leafhoppers, leafminers (including serpentine leafminer), cabbage looper, mites (including spider mites), thrips and whiteflies. Apply 25 lbs A, 14 DTH.
- SQUASH** - Aphids, amylworm, cucumber beetles (including spotted and striped cucumber beetles), fleabeetles, leafhoppers (including serpentine leafminer), cabbage looper, squash bug, squash vine borer, spider mites (including 2-spotted spider mites), stink bug, squash vine borer, stink bug, thrips and thrips. Apply 25 lbs A, 15 DTH.
- TOMATOES** - Aphids. Apply 15-15 lbs A, 7 DTH.
- TURTLE BEAT** - Ant, aphid, including green peach aphid, grasshopper, crane fly larvae, crickets, darkling beetle, fleabeetle, grasshopper, ground squirrel, hornworm, including tobacco hornworm, leafhoppers, cutworm, cabbageworm and vegetable weevil. Apply 25 lbs A, 100 sq yds. For crane fly larvae, flea beetle and thrips, apply 1-4 times as needed. For grasshopper and crane fly larvae, apply to burrowed areas of soil.
- TUBACOS** - Aphids, budworms, crane fly larvae, crickets, darkling ground beetle, flea beetle, grasshopper, hornworm, including tobacco hornworm, leafhopper, leafminer, meadow sparrow worm, stink bug, suckler, thrips and vegetable weevil. Apply 25 lbs A, 5 DTH for crickets, 15 DTH for thrips. Avoid contact with plant tissues when pinching or cutting tobacco. When necessary, re-enter treated fields within 24 hours after application. Protective clothing should be worn.
- TUMARIES** - Aphids, amylworm, Colorado potato beetle, fleabeetle, grasshopper, hornworm, including ground squirrel, leafhopper, leafminer (including serpentine leafminer), spider mites (including 2-spotted spider mites), stink bug, thrips, and thrips. Apply 25 lbs A, 10 DTH.
- TURNIPS** - Aphids, amylworm, cutworms (including diamondback moth), squash vine borer, spider mites, imported cabbageworm, cabbage looper, thrips, vegetable weevil and cabbage webworm. Apply 25 lbs A, 10 DTH.

## WARNING

Continued from front panel-

areas should not be re-entered until drifting insecticides and volatile residue has dissipated. Do not dust plants during pollination and/or flowering periods.

**FIRST AID TREATMENT:** Call a doctor (note for doctor): Therapeutic doses of atropine are effective as an antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine. **INTERNAL:** If swallowed make patient vomit by giving warm salt water or soapy water. Administer milk or water freely. Have patient lie down and keep warm. Never give anything by mouth to an unconscious patient. **SKIN:** Immediately remove all contaminated clothing and flush skin with water for at least fifteen minutes. **EYES:** Immediately flush with water for at least fifteen minutes. **INHALATION:** Give two atropine tablets at once if patient shows symptoms of blurred vision or pin point pupils.

Not for use or storage in or around the home

Toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Birds feeding on treated areas may be killed. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse containers. Destroy by burning in a safe place away from water supply.