

E-Z-Flo Diazinon[®] 3 Dust

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENT:

* O,O-Diethyl O-(2-isopropyl-6-Methyl-4-pyrimidinyl) phosphorothioate 3.0%

INERT INGREDIENTS:

97.0%

Total 100.0%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Harmful if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and food-stuffs. Avoid contact with clothing. Change contaminated clothing daily.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

Apply when insects first appear and repeat applications as infestations as infestations warrant. Generally, applications at 7-10 day intervals will be required. In many cases, it is advisable to dust field margins as well.

BROCCOLI AND CAULIFLOWER: Aphids, Imported Cabbage Worms, Diamond-back Moths - apply 20-50 lbs. per acre. Do not apply within 7 days of harvest.

CABBAGE: Aphids, Imported Cabbage Worms, Diamond-back moths - apply 20-50 lbs. per acre. Do not apply within 7 days of harvest.

MELONS, WATERMELONS AND SQUASH: Aphids, Leaf-hoppers, Serpentine Leaf Miners, Mites and Thrips - apply 20-45 lbs. per acre. Do not apply within 3 days of harvest on melons, winter squash, or watermelons nor within 7 days of harvest on summer squash.

CELERY: Aphids, Flea Beetles - apply 16.5 lbs. per acre. Do not apply within 10 days of harvest.

CUCUMBERS: Aphids - apply 20-50 lbs. per acre. Do not apply within 7 days of harvest.

LETTUCE: Aphids, Dipterous Leaf Miners - apply 16.5 pounds per acre. Do not apply within 10 days of harvest.

ONIONS: Onion Thrips - apply 20-40 lbs. per acre. Onion Maggots - apply 35-40 lbs. per acre. Do not apply within 10 days of harvest.

PEAS: Aphids - apply 16.5 lbs. per acre. Peas may be picked immediately. Peavines and peavine hay may be fed to dairy cattle, beef cattle, and sheep. Wait 4 days following treatment if forage is to be cut for hay.

PEPPERS: Aphids - apply 35-50 lbs. per acre. Do not apply within 5 days of harvest.

SNAP BEANS: Black Bean Aphids, Mexican Bean Beetles - apply 40-46 lbs. per acre. Do not apply within 7 days of harvest.

SPINACH: Aphids, Dipterous Leaf Miners - apply 16.5 lbs. per acre. Do not apply within 10 days of harvest.

STRAWBERRIES: Aphids, Cyclamen Mites, Two-spotted Mites - apply 20-40 lbs. per acre. Dust should be directed into the crowns of the plants and the plants sufficiently agitated to insure thorough coverage. Do not apply within 5 days of harvest.

TOMATOES: Aphids, Dipterous Leaf Miners, Drosophila flies - apply 20-40 lbs. per acre. Do not apply within 7 days of harvest.