

E-Z FLO THIODAN 3 DUST

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENT:

**Endosulfan (Hexachlorohexahydromethano-
2,4,3-benzodioxathiepin oxide) 3.00%**

INERT INGREDIENTS: 97.0%
Total 100.00%

*Thiodan is a registered trademark of Canadian Hoechst Limited.
U. S. Patent No. 2,799,685.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

BEANS (Succulent type): Use on Lima beans only when crop is to be harvested by combine. Do not use on Lima beans that are to be removed from the field for processing. — Black bean aphid, bean leaf skeletonizer, cucumber beetle, green stink bug, Mexican bean beetle. Use 17 to 33 lbs. per acre. Do not make more than 3 applications. Do not use within 3 days of harvest. Do not feed treated threshings to livestock or allow livestock to graze in treated fields.

BROCCOLI, CABBAGE: Cabbage looper, imported cabbage worm, and diamond back moth larvae. — Use 25 lbs. per acre. Apply when first signs of worm feeding appear and repeat as necessary. Do not apply within 7 days of harvest on Broccoli and cabbage.

BRUSSELS SPROUTS, CAULIFLOWER: Cabbage aphid, cabbage looper, cross striped cabbage worm, diamond back moth larvae, flea beetles, harlequin bug, imported cabbage worm. Use 25 to 33 lbs. per acre. Do not apply within 14 days of harvest. Unlimited number of applications.

CARROTS: Green peach aphid. Use 17 to 33 lbs. per acre. Do not make more than 2 applications. Do not apply within 7 days of harvest. Do not use tops for feed or feed.

CELERY (whole plant): Green peach aphid. Use 16 to 33 lbs. per acre. Do not apply within 7 days of harvest. Unlimited number of applications. Cabbage looper. Use 16 to 33 lbs. per acre. Do not make more than 3 applications. Do not apply within 4 days of harvest.

COLLARDS: Cabbage looper, diamond back moth larvae, harlequin bug, imported cabbage worm. — Use 25 lbs. per acre. Do not apply more than once. Do not apply within 21 days of harvest.

SWEET CORN (for fresh vegetable use only): Corn leaf aphid. Use 33 lbs. per acre. Apply when aphids first appear and repeat as required. Corn earworm. Use 12 lbs. per acre. Apply at 2 to 3 day intervals beginning when silks first appear and continuing until they begin to dry. Do not make more than 5 applications. May be applied up to day of harvest. Do not apply to sweet corn to be processed. Do not feed treated forage to any type of livestock or allow to graze in treated fields.

CUCUMBERS, MELONS, SUMMER SQUASH, WINTER SQUASH AND PUMPKINS: Aphids. Use 25 to 30 lbs. per acre. Apply when aphids first appear and repeat as required. Treat both upper and lower leaf surfaces thoroughly. No time limitations.

PEPPERS: Green peach aphid, flea beetles, hornworms, pepper maggot. — Use 33 to 50 lbs. per acre. Do not apply within 1 day of harvest. Apply when insects first appear and repeat as required.

ORNAMENTALS: Aphids. — Dust lightly, but uniformly. Apply when insects first appear and repeat as required. On *chrysanthemums* best results will be obtained if applied before plants flower.

EGGPLANTS: Green peach aphid, flea beetle, whitefly and Colorado potato beetle. — Use 20 lbs. per acre. Dust plants thoroughly, particularly the undersides of the leaves. Do not apply within 1 day of harvest.

POTATOES: Colorado potato beetle, potato flea beetle, potato leafhopper, aphids, southern armyworm, green stink bug, leaf-footed plant bug and potato tuberworm. — Use 30 to 33 lbs. per acre. Apply when insects first appear and repeat as required. No time limitations. Do not plant root crops other than carrots, potatoes, and sweetpotatoes as follow-up crops.

SPINACH: Green peach aphid, crown mite. — Use 25 lbs. per acre. Do not apply more than once. Do not apply within 21 days of harvest.

TOMATOES: Green stink bug, green peach aphid, blister beetle, Colorado potato beetle, flea beetle. — Use 20 lbs. per acre. Apply when insects first appear and repeat as required. Dust plants thoroughly, particularly the undersides of the leaves. Do not apply within one day of harvest.

TOMATOES: Whitefly. — Dust plants thoroughly. Apply to tomato plant seed beds in the field when whiteflies are first observed. Repeat as necessary to keep infestation as a low level. After seedlings are transplanted repeat applications as needed. Under field conditions after transplanting, apply at the rate of 15 to 20 lbs. per acre, depending upon the size of the plants. Do not apply within one day of harvest.

*Net Contents and Manufacturer
On Container*

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Avoid contamination of feed or foodstuffs. In case of contact with skin or eyes, flush with plenty of water. For eyes, get medical attention. Wash with soap and water after handling and before eating or smoking, wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas the day of application should wear protective clothing.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning, emptying, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.