

E-Z-FLO CAPTAN 5 DUST

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide) 5.00%

INERT INGREDIENTS: 95.00%

Total 100.00%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Avoid inhaling dust. Avoid contact with skin. Avoid contamination of feed and foodstuffs.

Consult local agricultural authorities for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated.

DIRECTIONS FOR USE

For best results apply when air is calm. A good hand or power dusting machine should be used to secure complete coverage of plants with a light film of dust. Begin application prior to or at first sign of infection. Repeat applications as necessary. Keep new growth protected.

RECOMMENDATIONS

APPLE: Scab, Frogeye Leaf Spot—40 to 70 lb. per acre as required to maintain adequate coverage in the pre-bloom and blossom applications.

CHERRY: Brown Rot, Leaf Spot—40 to 70 lb. per acre in the blossom and pre-harvest applications.

PEACH: Brown Rot and Scab—40 to 70 lb. per acre in the blossom and pre-harvest applications.

STRAWBERRIES: Botrytis Rot—45 to 60 lb. per acre beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking if conditions favor fruit rot.

BEANS, PEAS: Soil Treatment—Damp-off (Beans only), Root Rot—Work into top 3 to 4 inches of soil before planting. Apply 80 lb. per acre.

CELERY (Plant bed only)—Damp-off, Pink Rot—Apply 60 to 70 lb per acre when all seeds have germinated. Repeat if necessary.

CUCURBITS (Cantaloupe, Cucumber, Squash): Angular Leaf Spot, Anthracnose, Downy Mildew — Start applications when runners begin to form or at first sign of disease. Apply at rate of 30 to 40 lb. per acre on young plants increasing to 40 to 50 lb. per acre on mature plants.

POTATOES: Early and Late Blight—Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens. Apply 45 to 50 lbs. per acre on young plants. Use 60 lb. per acre on mature plants.

POTATO SEED PIECE TREATMENT: Damp-off, Seed Rot—Use at rate of 20 oz. per bushel cut seed pieces (32 oz. or 2 lb. to 20 lb. of cut seed pieces). Apply in convenient container or by dust attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates two days before bagging.

TOMATOES: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot—Apply 45 to 60 lb. per acre at weekly intervals on young plants; 60 to 70 lb. per acre on mature plants.

SOIL AND GREENHOUSE BENCH TREATMENT: 3 lb. per 1,000 sq. ft. (120 lb. per acre). Work into upper 3 or 4 inches of soil before planting. For damp-off and root rot diseases of seedlings and transplants of roses, other shrubs, flowers, lawn seedbeds, and vegetables such as beans, peas, celery, cabbage, eggplant, pepper and tomato.

ORNAMENTALS: Rose, Black Spot; Carnation, Rust Leaf Spot; Chrysanthemum, Flower Blight—Apply thoroughly prior to or at first signs of disease. Repeat applications at 7 to 10 day intervals. Shorten interval during periods of frequent rain and heavy dews.

LAWNS: Turf: Brown Patch, Damp-off, Leaf Spot, Root Rot—Apply 4 to 6 lbs, per 1,000 sq. ft. Repeat at 7 to 10 day intervals as needed. Melting out—Apply 4 to 6 lb. per 1,000 sq. ft., beginning when grass resumes growth in the Spring (March-April). Repeat at 7 to 14 day intervals as long as cool, wet weather persists.

Net Contents and Manufacturer
On Container