

E-Z-FLO SEVIN* 2 DUST

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate) 2%

INERT INGREDIENTS

98%

Total 100%

DIRECTIONS FOR USE

Apply when insects or signs of damage appear. Repeat 7 to 10 day intervals or as necessary unless indicated otherwise.

**ALFALFA, CLOVERS, COWPEAS (SOUTHERN OR FIELD PEAS), PEANUTS AND SOY-
BEANS:** For control of blister beetles and Mexican bean beetle, use 25 to 50 pounds per acre. For control of thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetles, Japanese beetle, alfalfa caterpillar, and Velvet Bean caterpillar use 50 pounds per acre. For control of armyworms, webworms, corn earworm, and stink bugs use 50 to 75 pounds per acre. Some leaf injury may occur if applications are made when tender foliage is wet.

BEANS: for control of Mexican bean beetle apply at the rate of 25 pounds per acre. For control of leafhoppers, bean leaf beetle, cucumber beetle, western bean cutworm, flea beetle and Japanese beetle apply at the rate of 50 pounds per acre. For control of corn earworm, armyworms and velvet bean caterpillar apply at the rate of 50 to 75 pounds per acre. For control of lima bean pod borer, Lygus and stink bugs apply at the rate of 100 pounds per acre.

CABBAGE, BROCCOLI, BRUSSEL SPROUTS, CAULIFLOWER AND KOHLRABI: For control of flea beetle and harlequin bug, apply at the rate of 25 to 50 pounds per acre. For control of corn earworm, army worm and cabbage caterpillars, and small cabbage loopers in home gardens or in commercial plantings apply at the rate of 50 to 100 pounds per acre.

CARROTS: For control of flea beetles and leafhoppers apply at the rate of 25 to 50 pounds per acre. For control of six-spotted leafhoppers (Aster yellows vector) apply at the rate of 50 to 75 pounds per acre. For control of spittlebugs, lygus and stink bugs apply at the rate of 50 to 100 pounds per acre.

CORN: For control of corn earworm, European corn borer, fall armyworm, sap beetles, and Japanese beetle apply at the rate of 50 to 100 pounds per acre. For larvae in the whorl (budworm damage) and foliage feeders apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on the severity of infestation. Timing and good coverage are absolutely essential for effective control.

CUCUMBERS, MELONS, PUMPKINS AND SUMMER AND WINTER SQUASH: For control of pickleworm, melonworm, cucumber beetles, leafhoppers, flea beetles and squash bugs apply at the rate of 50 pounds per acre.

FORAGE GRASSES AND PASTURE: For control of thrips, armyworms and stink bugs use 50 to 75 pounds per acre. For optimum thrips control in grasses grown for seed, direct dust into boot.

GRASSES: Apply 25 to 75 pounds per acre as often as necessary. Apply 25 to 50 pounds per acre on small plants or sparse vegetation, wasteland, and borders. Apply 50 to 75 pounds when grasshoppers are present.

LITTLE HEADS: For control of flea beetle, leafhoppers, harlequin bug apply at the rate of 25 to 50 pounds per acre. For control of six-spotted leafhopper apply at the rate of 50 to 75 pounds per acre. For control of corn earworm, armyworms, cabbage caterpillars, spittle bugs, Lygus and stink bugs apply at the rate of 50 to 100 pounds per acre.

OKRA: For control of corn earworm and stink bugs use 50 to 100 pounds per acre. Repeat at 5 to 7 day intervals or as necessary.

POTATO, TOMATO, EGGPLANT AND PEPPER: For control of Colorado potato beetle, flea beetles and leafhoppers apply at the rate of 25 to 50 pounds per acre. For control of tomato fruitworm, fall armyworms, tomato hornworms, European corn borer, lygus bugs and stink bugs apply at the rate of 50 to 100 pounds per acre. Apply as a foliage application only on potato.

SORGHUMS (including MILO AND SORGHUM): For control of armyworms, webworms, corn earworm and stink bugs use 50 to 100 pounds per acre. For optimum insect control on grain sorghum, direct dust into forming heads.

SUGAR BEETS: For control of leafhoppers, armyworms and webworms use 50 to 100 pounds per acre.

TOBACCO: In plant beds. For tobacco flea beetle, apply 2 1/2 to 5 pounds per 100 square yards when insects are present. Avoid excessive application. Plant injury may result. Repeat as needed at 10 to 14 day intervals and just before transplanting.

Injury may result when used on small plants or with application of heavy dosages on tender plants during rain or with heavy dew or fog. High for several days following application of dust. When applying, be sure to make sure not to overdose. Rotary dusters are recommended.

For control of tobacco flea beetle, apply 2 1/2 to 5 pounds per 100 square yards when insects are present. Avoid excessive application. Plant injury may result. Repeat as needed at 10 to 14 day intervals and just before transplanting.

*Trade Mark of Union Carbide Corporation for carbaryl insecticide. U. S. Patent Nos. 2,903,478; 3,009,855.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe Dust. Do not take internally. Avoid contact with skin. SKIN CONTACT MAY BE HARMFUL. Wash Hands Before Eating. Take Shower or Bath After Work. WEAR REGULAR LONG SLEEVED WORK CLOTHING. Change to Clean Clothing Daily. Do not store near feed and food products.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, Cholinesterase inhibitor. Atropine is antidotal.

PREHARVEST AND GRAZING USE LIMITATIONS: Do not use on crops to be harvested or grazed within 30 days of application. Do not use on crops to be harvested or grazed within 30 days of application. Do not use on crops to be harvested or grazed within 30 days of application.

Do not use on crops to be harvested or grazed within 30 days of application. Do not use on crops to be harvested or grazed within 30 days of application.

Net Contents and Manufacturer
On Container

