

THIRAM FUNGICIDE FOR THE CONTROL OF CERTAIN PLANT DISEASES

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Thiram (Tetramethylthiuram disulfide)...

INERT INGREDIENTS:

Contains DuPont "Thylate"* Thiram Fungicide.

*Registered Trademark and Product of E. I. duPont deNemours and Company, Incorporated.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

APPLES - For the control of Black Rot (fruit and foliage), Fly Speck, Sooty Blotch, Bitter Rot, Brooks Spot, Apple Blotch, Botry -- Osphaeria Spot and Black Pox - Apply 35 to 40 pounds per acre during pre-bloom, calyx, and early cover dustings. Whenever weather conditions favor disease, shorten the interval between applications. 5% Thylate Dust is a protective fungicide;

PEACHES — Brown Rot — For blossom blight, apply at 3 to 4 day intervals during bloom at a rate of 40 pounds per acre, timed to keep newly opened blossoms covered. For fruit rot, apply at petal fall and 2 weeks later. Make one or more additional applications at 7 day intervals beginning about one month before harvest, or earlier if rot is apparent on green fruit.

Rhizopus Rot - Apply as directed for fruit phase of brown rot.

Scab – Apply at petal fall at the rate of 40 pounds per acre, and repeat 3 to 6 times at 10 to 14 day intervals or until disease is well under control. Do not apply within 7 days of harvest.

STRAWBERRIES -- For control of gray mold (Botrytis) - Use at the rate of 40 lbs. per acre. Make 3 to 5 applications at ten-day intervals, beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results.

If applied within 3 days of picking, remove residues by washing.

May cause irritation of eyes, nose, throat, and skin. Harmful if inhaled or swallowed. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. In case of contact, wash skin with soap and water; for eyes, flush with plenty of water and get medical attention. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

53-263-660

611-1R

hence, applications are most effective when made in advance of infection periods.

Good coverage of foliage and developing fruit is required for best results.

CAUTION

U.S.D.A. Reg. No. 635-218