


E-Z-Flo Copper Zinc Chromate Thiodan® 6-3 Dust



CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Endosulfan (hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide)	3.00%
Copper-zinc-chromate complex* **	5.70%
INERT INGREDIENTS:	91.30%
Total	100.00%

* 15 CuO-10ZnO-6 CrO3-25 H₂O (licensed under U.S. Patent 2,573,738)

** Active Ingredients expressed in terms of metal content: Copper, 1.8%; Chromium, 0.6%; Zinc, 1.2% (Total 3.6)

* Thiodan is a registered trade mark of Canadian Hoechst Limited. U. S. Patent No. 2,799,685.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds, and streams. Do not apply in any manner not specified on the label. Do not apply when weather conditions favor drift of dust from crop areas treated.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

DIRECTIONS FOR USE

This dust is formulated to control certain insects and diseases on potatoes. In general, applications should be made at 7 to 14 day intervals, depending on weather conditions and the severity of insect or disease problems. For disease control, applications should be started before diseases occur.

Local weather and other conditions may affect the use of this material. State agricultural experiment stations may issue recommendations to fit local conditions.

POTATOES: For control of aphids, Colorado Potato Beetle, Leaf-hopper and Flea Beetle, Southern Armyworm, Green Stink Bug, Leaf-footed Plant bugs, Potato Tuberworms, and Whitefly; for the Control of Early and Late Blight of Potatoes: Apply this dust at the rate of 30 to 33 pounds per acre at 7 to 14 day intervals, depending on weather conditions and insect and disease incidence. Under conditions which favor Blight development, more frequent applications are required than during dry, hot weather. Begin dusting soon after potato plants show above ground; continue throughout the growing season. Thorough coverage is important. No time limitations. Do not plant root crops other than carrots, potatoes, and sweet potatoes as follow-up crops.

TOMATOES: For control of Aphids, Colorado Potato Beetle, Flea Beetle, Whitefly, Green Stink Bug, Blister Beetle, Early Blight and Late Blight. Apply at the rate of 33 pounds per acre. Do not apply within 1 day of harvest.

CUCUMBERS: For the control of Downy Mildew, Anthracnose, Scab, Mucrosporium Leaf Spot, Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle and the Squash Vine Borer. Apply at the rate of 33 pounds per acre when plants are two to six inches high and continue as needed.

MELONS: For the control of Downy Mildew, Anthracnose, Scab, Mucrosporium Leaf Spot, Aphids, Cucumber Beetle, Striped Cabbage Flea Beetle, Cabbage Looper, Omniverous Leafroller. Apply at the rate of 33 to 50 pounds per acre. Apply when plants are two to six inches high and continue as needed.

SUGAR BEETS: For control of Cercospora Leaf Spot, Green Peach Aphid and the Sugar Beet Webworm. Apply at the rate of 25 to 33 pounds per acre. Apply at 10 day intervals as needed. Do not feed treated tops to livestock.

Net Contents and Manufacturer
On Container