

ACCEPTED

June 13, 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRA-
ED UNDER NO. 635-245



E-Z FLO

CAPTAN-STREPTOMYCIN

7.5-.01

POTATO SEED PIECE PROTECTANT

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN

Net Contents and Manufacturer
— On Container

ACTIVE INGREDIENTS:

*Captan 7.50%

**Streptomycin01%

INERT INGREDIENTS: 92.49%

100.00%

*N-trichloromethyl thio-4-cyclohexene-1,2-dicarboximide

**Streptomycin from streptomycin sulfate (Agricultural grade)

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

This material is especially formulated for potato seed piece treatment.

POTATO (Seed Piece Treatment): For use as an aid in the control of Tuber Soft Rot, Black Leg and surface borne Verticillium Wilt. Dust at 10 ounces per bushel cut seed pieces (16 ounces or 1 lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 to 2 days, the treated pieces should be stored for 2 or 3 days in open crates before bagging.

Do not use treated seed for food or feed purposes.

CAUTION

This product may cause skin irritation. Avoid contact with skin and eyes. Avoid breathing dust. Wash with soap and water after exposure. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

STORE IN COOL DRY PLACE — PROTECT FROM EXCESSIVE HEAT.

Not licensed for use or export outside of the United States, its Territories and Possessions, Canada or Mexico.

U. S. Pat. Nos. 2,553,770 — 2,553,771.