

# E-Z FLO SEVIN\* 10 DUST

**CAUTION:**  
**KEEP OUT OF THE REACH OF CHILDREN**

ACCEPTED  
MAY 6 1978  
635-252  
U.S. DEPT. OF AGRICULTURE  
GENERAL INSECTICIDE  
ACT  
POISON REGISTER-  
NO.

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl  
N-methylcarbamate)\* ..... 10.0% by wt.

**INERT INGREDIENTS:** ..... 90.0% by wt.  
100.0%

\*Trade Mark of Union Carbide Corporation for carbaryl insecticides.  
U. S. Pat. No. 2,903,478

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

## DIRECTIONS FOR USE

**FOR CONTROL FOR MAJOR TOBACCO INSECTS:** In the field, For Hornworms, Budworms, Aphids, Flea Beetles, Suckfly, Grasshoppers and Japanese Beetles, apply 15 to 20 lbs. per acre depending on the size of the plants. Make applications when insects first appear and repeat as necessary. Avoid excessive applications.  
**FOR BUDWORMS:** When hand dust applications are directed to the bud and top leaves only, apply 4 to 6 lbs. per acre. Avoid heavy applications with cloth bags as plant injury may result. Do not apply within 3 days of priming or cutting. Avoid getting wet with plant juices when cutting the crop. Injury may occur on tender foliage in the presence of rain or high humidity of several days duration following application of Sevin.

### DIRECTIONS FOR USE FOR ORCHARD DUSTING

**ON APPLES AND PEARS:** For control of codling moth, green apple aphid, apple maggot, red-banded leaf roller, pear psylla and periodical cicada.

**ON PEACHES:** For control of Oriental fruit moth, peach twig borer and periodical cicada.

**ON CHERRIES, PLUMS AND PRUNES:** For control of cherry fruit fly, codling moth, Eastern tent caterpillar, fruit tree leaf roller, peach twig borer, orange tortrix, plum curculio, red-banded leaf roller.

Use 40-60 pounds of Sevin 10% Dust per acre. Repeat applications at 10-day intervals if needed. SEVIN is toxic to honeybees; avoid use during flowering and pollination period. Application of SEVIN within 30 days after full bloom may also provide fruit thinning; to avoid this, delay use until at least 30 days after bloom. 10% Dust may be applied up to one day before harvest.

**APRICOTS AND NECTARINES:** For control of Oriental fruit moth, peach twig borer, codling moth, orange tortrix, tussock moth, fruit tree leaf roller, Japanese beetle and periodical cicada, use 40 to 60 lbs. per acre. Repeat applications at 10 day intervals if needed. Do not apply within 3 days of harvest.

**CRANBERRIES:** For control of fruitworms, fireworms, leafhoppers and cutworms use 20 to 40 lbs. per acre. Apply at late bloom and repeat as needed at 7 to 10 day intervals. Do not apply within one day of harvest.

### VEGETABLE, VINE AND FIELD CROPS:

**COTTON: SEEDLING STAGE:** Thrips — Use 8 lbs. per acre for 2-3 applications at 7 day intervals when first 2 leaves begin to spread. **BEGINNING OF SQUARING:** Overwintered Boll Weevils, Flea Hoppers, Plant Bugs, Boll Worm — Use 10 lbs. per acre for 3 applications at 5-7 day intervals. **MAJOR FRUITING SEASON:** Boll Weevil, Boll Worm, Cotton Leaf Worm, Lygus Bug, Cotton Leaf Perforator — Use 10-20 lbs. per acre at 4-5 day intervals at first sign of insects. Continue application until top bolls are fully grown.

**BEANS:** Mexican bean beetle, Leafhoppers — Apply 10 lbs. per acre. Lygus and Stink bugs — Apply 20 lbs. Apply thoroughly to upper and lower leaf surfaces.

**CORN: Corn Earworm** — Apply 15-20 lbs. per acre starting when the first silks appear. Repeat at 2-3 day intervals throughout the silking period. Direct duster outlets toward ear zone to insure thorough coverage of each silk. May be applied up to harvest on sweet corn ears for human consumption.

**CUCUMBER, SUMMER SQUASH:** Flea beetle, melonworm, pickleworm, spotted cucumber beetle, squash bugs — Apply 10 lbs. per acre.

**GRAPES:** Grape leaf roller — Apply 10-15 lbs. per acre just before first brood larvae emerge from the first rolls. Apply 10-20 lbs. for the control of 2nd and 3rd broods. Grape leafhopper — Apply 10 lbs. May be expected to control DDT-resistant leafhoppers.

**EGGPLANT, PEPPER, TOMATO** — Tomato fruitworm, fall armyworm, flea beetles — Apply 10-20 lbs. per acre. European corn borer, stink bugs, tomato hornworm — Apply 20-40 lbs.

**POTATO:** Colorado potato beetle, leaf hopper, flea beetle and Lygus bugs — Apply 20 lbs. Apply thoroughly to upper and lower leaf surfaces.

**MAY BE APPLIED UP TO HARVEST ON THE ABOVE VEGETABLE, VINE AND FIELD CROPS.**

**STRAWBERRIES:** Meadow spittlebug, strawberry leaf roller — Apply 20 lbs. per acre. May be applied up to 1 day before harvest.

Consult state agricultural experiment stations for additional information as the timing, number, and rate of applications needed will vary with local conditions.

## CAUTION

Harmful if inhaled or swallowed. Do not take internally. Avoid prolonged breathing of dust. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeve work clothing. Change to clean clothing daily.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Net Contents and Instructions on Container