

(Contains 6 Lbs. Toxaphene Per Gallon)

WARNING: **KEEP OUT OF THE REACH OF CHILDREN**

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Toxaphene*	59.0%
Xylene	
INERT INGREDIENTS:	3.0%
-	

*Technical Chlorinated Camphene (Chlorine Content 67% to 69%). .

The manufacturer makes no warranties, express or implied, concerning this product for its use, which extend be-yond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY **E-Z FLO CHEMICAL COMPANY DIVISION OF KIRSTO COMPANY**

COLUMBUS, OHIO

LANSING, MICHIGAN

11-25

53-263-846

U.S.D.A. Reg. No. 635-263

DIRECTIONS FOR USE

For dilution with water to form emulsions suitable for control of GRASSHOPPERS. ARMYWORMS, AND GREEN CUTWORMS. Application may be made in 3 to 10 gals. of water by aerial equipment or in 10 to 150 gals, of water by conventional or low volume ground equipment per recommended practices to give thorough coverage. The following suggestions are general recommendations. For timing, number of applications and specific recommendations consistent with label directions, consult your State agricultural experiment station

GRASSHOPPERS: Barley, corn, oats, rye and wheat — Use 1 to 1¹/₃ quarts of this concentrate (1¹/₂ to Z lbs. Technical Toxaphene) with sufficient water to give adequate coverage per acre. For best results, treatment should be made before grasshoppers reach the flying stage. More than one treatment may be necessary.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

ARMYWORMS (Fall, Southern or Yellow Striped) and GREEN CUTWORMS: Barley, corn, oats, rye and wheat - Use 11/2 to 42/3 pts. per acre in sufficient water for good coverage.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

BARLEY, OATS, RYE AND WHEAT: Fall Armyworm, Southern Armyworm, or Yellow Striped Armyworm -- Use 11/3 to 21/3 qts. per acre in sufficient water for coverage.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

COTTON INSECTS:

EARLY SEASON: Thrips and Cutworms Control — If thrips are a problem during moist years make 2 or 3 applications using 34 to 1 qt. per acre at weekly intervals. Use 11/2 to 21/2 gts. per acre for Cutworm Control directed at base of plants when worms appear. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

100.0%

WARNING

May be fatal if swallowed. Do not breathe spray mist. Do not get on skin, in eyes, or on clothing. Wash thoroughly after using. Do not store near feed and food products.

Do not use this product on dairy animals or in dairy barns. Do not feed treated forage to dairy animals or animals being finished for slaughter.

Highly toxic to fish and wildlife. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply under conditions favoring drift outside the target area. Birds feeding on treated areas may be killed.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

DESTROY CONTAINER WHEN EMPTY NEVER RE-USE.

Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).