E-Z FLO ROTENONE 0.75 D

CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS: Rotenone	0.75%
Other Cube Resins	1.50%
INERT INGREDIENTS:	97.75%
Total	100.00%

CAUTION

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Destroy Container When Empty. Never Re-use.

51**-2**9

EPA Est. 635-MI-1; OH-1

E.P.A.Reg. No. 635-269

DIRECTIONS FOR USE

Control. Always repeat dusting after a rain. Do not apply within I day of harvest.

BEANS: Black bean aphid, cucumber beetle, Mexican bean beetle- Use 37' lbs. per acre.

BROCCOLI, CABBAGE: Cabbage looper, imported cabbageworm. Use 183 lbs. per acre. Apply when first signs of worm feeding appear and repeat as necessary.

BRUSSELS SPROUTS, CAULIFLOWER: Cabbage aphid, cabbage looper, cross-striped cabbageworm, flea beetles, young harlequin bug, imported cabbageworm-Use 18\frac{3}{4} lbs. per acre.

CARROTS: Green peach aphid- Use 183 lbs. per acre.

CELERY: Green peach aphid- Use 183 lbs. per acre.

COLLARDS: Cabbage looper, young harlequin bug, imported cabbageworm- Use 18³/₄ lbs. per acre.

CUCUMBERS, MELONS, SUMMER SQUASH, WINTER SQUASH and PUMPKINS: Aphids – Use 18\frac{3}{4} lbs. per acre. Apply when aphids first appear and repeat as required. Treat both upper and lower leaf surfaces thoroughly.

PEPPERS: Green peach aphid - Use 18% lbs. per acre. Apply when insects first appear and repeat as required.

EGGPLANTS: Green peach aphid, flea beetle, and Colorado potato beetle-Use 37½ lbs. per acre. Dust plants thoroughly, par'icularly the undersides of the leaves.

POTATOES: Colorado potato beetle, aphids - Use 37^{1}_{2} lbs. per acre. Apply when insects first appear and repeat as required.

SPINACH: Green peach aphid- Use 183 lbs. per acre.

TOMATOES: Green peach aphid, Colorado potato beetle, flea beetle- Use 37°_2 ibs. per acre. Apply when insects first appear and repeat as required. Dust plants thoroughly, particularly the undersides of the leaves.