

DIRECTIONS FOR USE

TOBACCO - PLANT BEDS: Blue Mold Control - Start dusting when plants are about the size of a dime, or when disease is first reported in area. Dust at the rate of 2 to 4 pounds per 100 square yards. (100 foot x 9 foot bed.) Apply at 3 to 4 day intervals until transplanting time. Always repeat dusting after


CAUTION

May cause irritation of eyes, nose and throat and skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water; for eyes, get medical attention.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of

Container disposal: Do not re-use container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

U.S.D.A. Reg. No. 635-275



11-22

53-263 647

DIRECTIONS FOR USE

TOBACCO - PLANT BEDS: Blue Mold Control - Start dusting when plants are about the size of a dime, or when disease is first reported in area. Dust at the rate of 2 to 4 pounds per 100 square yards. (100 foot x 9 foot bed.) Apply at 3 to 4 day intervals until transplanting time. Always repeat dusting after

CAUTION

May cause irritation of eyes, nose and throat and skin. Avoid breathing dust. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water; for eyes, get medical attention.

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of

Container disposal: Do not re-use container Destroy by burying with waste or burning. Stay away from smoke or fumes.

USDA Reg No 635-275