

E-Z FLO CHLORO 8 SPRAY

WARNING

KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Chlordane*, Technical 72.0%
Xylene 23.0%

INERT INGREDIENTS: 5.0%
100.0%

*Equivalent to 43.2% Octachloro-4, 7-Methanotetrahydroindane and 28.8% related compounds.

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY
E-Z FLO CHEMICAL COMPANY
DIVISION OF KIRSTO COMPANY
COLUMBUS, OHIO LANSING, MICHIGAN

15-19

53-263-887

E.P.A. Reg. No. 635-281

DIRECTIONS FOR USE

DIRECTIONS FOR USE: Pour the required amount of this material through screen into partially filled spray tank with agitator running. Keep agitator running while mixing and spraying. Mixture does not mix well readily, but tends to separate as an oily layer. Do not use as injury to plants may result.

CORN (field and sweet)

Preplant Broadcast application: Make broadcast application of 2.5-4 pints per acre in 15 to 20 gallons of water per acre, for control of the soil insect complex, including wireworms, white grubs, seed corn maggot, and larvae of green June bug, Asiatic garden beetle, Japanese beetle, and white grub beetles. For the control of corn rootworms (Western, Southern, Northern), cutworms, cornfield ants, and corn root aphids, apply 2-4 pints per acre.

Apply after the field is plowed. Disk in promptly. Use the higher rate on heavier soils and the lower rate on light sandy soils. Use the higher rates for heavy infestation of cutworms.

Row application: To control corn rootworms, (Western, Northern, Southern), cornfield ants and corn root aphids, apply 1 pint per acre at planting time in the furrow, behind the planter shoe and ahead of the press wheel. To control wireworms and white grubs, apply 2½ pints per acre in the row at planting time.

Consult your State Agriculture Extension Service for detailed recommendations.

LAWNS: Crabgrass — Apply before crabgrass seeds germinate in the early spring. Use 2 pints per 1,000 sq. ft. in enough water for thorough coverage. **CAUTION:** Do not graze dairy animals or animals being finished for slaughter on treated area. Avoid inhalation. Avoid contamination of feed and foodstuffs. Do not use on newly seeded lawns. Keep children and pets off treated area until this material has been thoroughly washed into the soil and the grass has completely dried.

LAWN INSECTS: Chiggers, ticks, crickets — Use 1½ tablespoonfuls in 5 gallons of water per 1,000 sq. ft. Do not sprinkle for 48 hours after application. **Ants, grasshoppers** — Use 1¼ tablespoonfuls in 5 gallons of water per 1,000 sq. ft. Do not sprinkle for 48 hours after application. **Webworms, chinch bugs, false chinch bugs, cutworms** — Use 3 tablespoonfuls in 5 gallons of water per 1,000 sq. ft. Sprinkle thoroughly after application. **White grubs** — Use 3½ tablespoonfuls in 5 gallons of water per 1,000 sq. ft. Sprinkle thoroughly after application. **Male crickets** — Use 1½ tablespoonfuls in 5 gallons of water per 1,000 sq. ft. Sprinkle well after application.

TERMITES: Use ½ gallon to 1 gallon of this material to 100 gallons of water and apply at the rate of 1 gallon per linear foot to a trench 1 to 2½ feet deep and 1 to 12 inches wide dug next to the foundation. Make trench 30 inches deep for buildings with deep foundations. Inspect annually for possible reinfestation with termites. Do not dig trench below the level of the footing.

ACCEPTED

OCT 6 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 635-281

WARNING

If swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTIDOTE: External — In case of spillage on skin, wash promptly with soap and water.

Internal — Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

NET CONTENTS ON CONTAINER

General Information

Avoid drift onto adjoining food and forage crops about to be harvested. Use only as recommended on label. Consult your State Agricultural Experiment Station for timing of applications.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE: Do not combine emulsifiables with wettable powders in the same spray tank since they may not be compatible.

COMBUSTIBLE MIXTURE — DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. PROTECT FROM TEMPERATURES BELOW 40° F.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.