

LIME SULFUR SOLUTION

CAUTION

KEEP OUT OF THE REACH OF CHILDREN
SEE SIDE OF PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:

Calcium Polysulfide	29%
INERT INGREDIENTS	71%
	<hr/>
	100%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Seller guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within lawful limits and (b) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states. Seller makes no other express or implied guarantee, warranty or representation. **No Warranty of Merchantability or of Fitness for a Particular Purpose is Implied.**

SOLD BY

E-Z FLO CHEMICAL COMPANY
DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

04-R

53-263-896

U. S. D. A. Reg. No. 635-292

DIRECTIONS FOR USE

DORMANT SPRAYING: Peaches — Peach leaf curl — Dilute E-Z-Flo Lime-sulfur solution in water at 1 to 15. Apply in late dormant period just before the buds begin to swell.

PEACHES — San Jose Scale — Dilute E-Z-Flo Lime-sulfur solution in water at 1 to 8. For dormant application only.

APPLES — San Jose Scale — Dilute E-Z-Flo Lime-sulfur solution in water at 1 to 8. Apply as a dormant spray only.

SUMMER SPRAYING: Apples — Apple scab and certain other fungus diseases — For delayed dormant, pre-pink, and pink applications, use 1½ to 2 gals. of E Z Flo Lime-sulfur solution to 100 gals. of water. Follow your local experiment station recommendations regarding usage for this purpose in your area.

ACCEPTED

MAY 23 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 635-292

CAUTION: May cause skin irritation—Avoid long continued or frequently repeated contact. In case of contact, wash thoroughly with plenty of water.

NET CONTENTS OF CONTAINER

Density - Baume at 60°F - 31°

Contains 3 lbs. per gal.
Calcium & Sulfur expressed
as gypsum and 1.9 lbs. per
gal. other combined sulfur.

Product weight - 10.59 lbs.
per gal. at 68°F.

ACTIVE INGREDIENTS:	
Calcium Polysulfide	29%
INERT INGREDIENTS	71%
	<hr/>
	100%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Seller guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within the limits and (b) that the directions, warnings and other statements on this label are based upon responsible expert evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states. Seller makes no other express or implied guarantee, warranty or representation. **No Warranty of Merchantability or of Fitness for a Particular Purpose is Implied.**

SOLD BY
E-Z FLO CHEMICAL COMPANY
DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

94-R

53-263-896

U. S. D. A. Reg. No. 635-292

DIRECTIONS FOR USE

DORMANT SPRAYING: Peaches — Peach leaf curl — Dilute E-Z-Flo Lime-sulfur solution in water at 1 to 15. Apply in late dormant period just before the buds begin to swell.

PEACHES — San Jose Scale — Dilute E-Z-Flo Lime-sulfur solution in water at 1 to 8. For dormant application only.

APPLES — San Jose Scale — Dilute E-Z-Flo Lime-sulfur solution in water at 1 to 8. Apply as a dormant spray only.

SUMMER SPRAYING: Apples — Apple scab and certain other fungus diseases — For delayed dormant, pre-pink, and pink applications, use 1½ to 2 gals. of E-Z-Flo Lime-sulfur solution to 100 gals. of water. Follow your local experiment station recommendations regarding usage for this purpose in your area.

NOTE: Do not apply Lime-sulfur when trees are in bloom, and do not apply after bloom within two weeks of any oil spray. Do not apply during periods of high temperatures, since sulfur sprays may burn foliage under such conditions. Do not apply this material during freezing weather. Use with care near buildings, since discoloration of painted surfaces may result.

3 teaspoons = 1 tablespoon	10 2/3 tablespoons = 2/3 cups	1 cup = 1/2 pt. = 8 oz.
4 tablespoons = 1/4 cup	12 tablespoons = 3/4 cups	2 cups = 1 pt. = 16 oz.
5 1/3 tablespoons = 1/3 cup	16 tablespoons = 1 cup	4 cups = 1 qt. = 32 oz.
8 tablespoons = 1/2 cup	2 tablespoons = 1 liquid ounce	4 qts. = 1 gal. = 128 oz.
2 qts. of any material per acre in 25 gal. of water = 5 tablespoonful in 1 gal. water		
1 qt. of any material per acre in 25 gal. of water = 2½ tablespoonful in 1 gal. water		
1 pt. of any material per acre in 25 gal. of water = 4 teaspoonful in 1 gal. water		
½ pt. of any material per acre in 25 gal. of water = 2 teaspoonful in 1 gal. water		

quently repeated contact. In case of contact, wash thoroughly with plenty of water.

NET CONTENTS OF CONTAINER

Density - Baume at 60°F - 31°

Contains 3 lbs. per gal.
 Calcium & Sulfur expressed as gypsum and 1.9 lbs. per gal. other combined sulfur.

Product weight - 10.59 lbs. per gal. at 68°F.