E-Z FLO METHYL PARATHION 2E

(Contains 2 Lbs. Methyl Parathion Per Gal.)

DANGER:
KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Additional Warnings and Antidote Statements

POISON

ACTIVE INGREDIENTS:

0,0-dimethyl 0-p-nitrophenyl thiophosphate	**24.9%**
Xylene	**68.8%**
INERT INGREDIENTS:	**6.3%**
	100.0%

Seller guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within lawful limits and (b) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all States. Seller makes no other express or implied guarantee, warranty, or representation. **No Warranty of Merchantability or of Fitness for a Particular Purpose is Implied.**

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA **COLUMBUS, OHIO** **LANSING, MICHIGAN**

94-24 53-263-870 U.S.D.A. Reg. No. 635-296

DIRECTIONS FOR USE

COTTON INSECTS: Aphids, Red Spiders, Cotton Leafworms, Flea Hoppers, Lygus Bugs and other plant bugs, Boll Weevil and Thrips. Workers entering fields within 24 hours after application should wear protective clothing.

SEEDLING STAGE: For Thrips control use ½ to 1 pint per acre when first 2 leaves spread; 2 or 3 applications at 7 day intervals.

BEGINNING OF SQUARING AND MAJOR FRUITING SEASON: When any of the above cotton insects appear use 1 to 2 pints per acre at 4 to 5 day intervals until insects are under control or until top bolls are fully grown. In case of heavy Boll Weevil migration, shorten intervals to 3 days. Do not apply within 5 days of hand picking.

ALFALFA: Two Spotted Aphid—Apply 1 to 1½ pts. per acre. Alfalfa Weevil Larvae and Adults — Apply 2 pts. per acre when buds and shoots first appear ragged or when larvae count reaches 25 per sweep. Workers entering fields within 24 hours after application should wear protective clothing. Do not apply within 15 days of harvest.

BARLEY, OATS, and WHEAT: For Spotted Alfalfa Aphid use 1 to 1½ pints per acre. For aphids on wheat, oats or barley use 1 to 1½ pints per acre. Do not apply within 15 days of harvest. Workers entering fields within 24 hours after application should wear protective clothing.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE

NET CONTENTS ON CONTAINER

WARNING

[illegible]

ANTIDOTE IF SWALLOWED: Give a tablespoonful of salt in a glass of warm water and induce vomiting until fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY.

ANTIDOTE IF ON SKIN: Wash immediately with soap and water.

NOTE TO PHYSICIAN: Methyl Parathion is a cholinesterase inhibitor and treatment of severe poisoning should include the use of large doses of atropine and artificial respiration if necessary. Observe patient continuously for 48 hours.

SYMPTOMS OF POISONING: A sense of tightness in the chest. Vomiting and diarrhea. Stomach pains. [illegible] persons but not operating areas [illegible] may be dangerous to drift. Treated areas should not be re-entered until drifting insecticide [illegible] have dissipated.

USE PRECAUTIONS

This product will kill wildlife. Apply only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use on sorghum. Do not apply or allow to drift where dairy animals or animals being finished for slaughter may be grazing. Do not apply within 5 days of hand picking. Workers entering fields within 24 hours after application should wear protective clothing.

Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning (keep out of smoke).

CAUTION: Do not use or store near heat or open flame.

ACCEPTED

Jun 3, 1969

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR LIMITED USE ON COTTON REGISTRATION NO. 635-296

E-Z FLO METHYL PARATHION 2E

(Contains 2 Lbs. Methyl Parathion Per Gal.)

DANGER:
KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Additional Warnings and Antidote Statements

POISON

ACTIVE INGREDIENTS:

0,0-dimethyl 0-p-nitrophenyl thiophosphate	24.9%
Xylene	68.8%
INERT INGREDIENTS:	6.3%
	100.0%

Seller guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within lawful limits and (b) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all States. Seller makes no other express or implied guarantee, warranty, or representation. **No Warranty of Merchantability or of Fitness for a Particular Purpose is Implied.**

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA **COLUMBUS, OHIO** **LANSING, MICHIGAN**

94-24 53-263-870 U.S.D.A. Reg. No. 635-298

DIRECTIONS FOR USE

COTTON INSECTS: Aphids, Red Spiders, Cotton Leafworms, Flea Hoppers, Lygus Bugs and other plant bugs, Boll Weevil and Thrips

WARNING

USE PRECAUTIONS

This product will kill wildlife. Apply only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from