

E-Z FLO ENDRIN-METHYL PARATHION 1.6-2 EMULSIVE
 (CONTAINS 1.6 LBS. ENDRIN AND 2 LBS. METHYL PARATHION PER GAL.)

DANGER:
KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Additional Warnings and Antidote Statements



POISON



ACTIVE INGREDIENTS:
 Endrin (Hexachlorocyclohexane-endo, endo-dimethanonaphthalene) 23.14%
 O,O-dimethyl O-p-nitrophenyl thiophosphate 53.93%
 Xylene range Aromatic Hydrocarbon Solvent 3.95%

INERT INGREDIENTS:

18.98%
 23.14%
 53.93%
 3.95%
100.00%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA
 COLUMBUS, OHIO

LANSING, MICHIGAN
 LEXINGTON, KENTUCKY

86-26

53-263-897

U.S.D.A. Reg. No. 635-315

DIRECTIONS FOR USE

Apply using conventional ground equipment using the specified dosage of material per acre in 50 to 150 gallons of water per acre to give complete coverage.

COTTON INSECTS:

SEEDLING STAGE: Aphids, Thrips, Fleahoppers - Use 1/2 pt. per acre for 2 or 3 applications at 7 day intervals beginning when first two leaves spread.

BEGINNING OF SQUARING: Overwintered Boll Weevils, Bollworms, Fleahoppers - Use 1 pint per acre for 3 applications at 7 day intervals.

MAJOR FRUITING SEASON: Boll Weevil, Bollworm, Cotton Leafworm, Cabbage Looper, Lygus Bug, Aphids, Plant Bugs and Red Spiders - Use 1 to 2 pints per acre at 4 to 5 days intervals at first sign of insects. Continue at present rate until bolls are fully grown.

Do not

APPLY WITHIN 5 DAYS OF HARVEST. Do not allow workers entering fields within 24 hours after application. Wear protective clothing.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DANGER

Rapidly absorbed through skin. Poisonous if swallowed, inhaled or absorbed through skin. Do not inhale, get in eyes, on hair or on clothing. Avoid contact with food. This type passed by the U.S. Department of Agriculture for Methyl Parathion and Endrin mixture protection. Wear protective clothing and goggles. In case of contact with skin immediately wash with soap and water. Remove all contaminated clothing. Flush skin or eyes with plenty of water for at least 15 minutes. In case of contact for eyes get medical attention. Do not contaminate feed or foodstuffs. Wash all contaminated clothing before reuse. Keep out of reach of children and domestic animals. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be reentered until drifting insecticide and volatile residues have dissipated.

SYMPTOMS OF POISONING: Irritation of the skin, vomiting, diarrhea, dizziness, headache.

ANTIDOTE IF SWALLOWED: Give 1-2 glasses of water with 1-2 teaspoons of activated charcoal. Do not induce vomiting. If ON SKIN: Wash immediately with soap and water.

NOTE TO PHYSICIAN: Methyl Parathion is a cholinesterase inhibitor. Symptoms of poisoning include: blurred vision, sweating, salivation, nausea, vomiting, diarrhea, muscle twitching, and respiratory distress. 2-PAM is also antidotal and may be administered in conjunction with atropine. Never give morphine.

2-PAM is also antidotal and may be administered in conjunction with atropine. Never give morphine.

Do not apply or allow to drift where dairy animals or animals being finished for slaughter may be grazing. Do not graze dairy animals or animals being finished for slaughter on treated fields. Workers entering fields within 7 days after treatment should be protected. Do not apply within 7 days of hand picking.

USE PRECAUTIONS

CAUTION: Do not use or store near heat or open flame.

CONTAINER DISPOSAL: Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.)

Do not apply or allow to drift where dairy animals or animals being finished for slaughter may be grazing. Do not graze dairy animals or animals being finished for slaughter on treated fields. Workers entering fields within 7 days after treatment should be protected. Do not apply within 7 days of hand picking.

This product will kill fish and wildlife. Keep out of any body of water. Do not apply when weather conditions favor drift of spray from areas treated. Do not apply where runoff is likely to occur. Birds feeding on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.