

E-Z FLO Phygon-Sulphur Dust

ACTIVE INGREDIENTS:	
Dichlone (2,3-Dichloro-1, 4-Naphthoquinone)	.50%
Sulphur	84.00%
INERT INGREDIENTS:	15.50%
TOTAL	<u>100.00%</u>

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION


Dichlone is a skin irritant, especially during warm weather. Irritation is similar to sunburn, evidence fails to show any permanent injury. Avoid inhalation of dust. Wear protective clothing. Wash hands and face thoroughly with soap and water after use, before smoking or eating. Keep away from feed or food products. Protective creams are suggested for the hands and face of sensitive individuals when dusting with Dichlone.

Do not use within 10 days of an oil spray. Sulphur may burn foliage when temperatures are excessively high.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

Net Contents and More
On Container



DIRECTIONS FOR USE

USE ONLY AS DIRECTED. Apply at the rate of 50 pounds per acre or 1/2 pound on each side for full grown apple and sweet cherry trees or 1/4 pound to peaches, plums and red tart cherries.

APPLES - For control of apple scab, begin applications at the first sign of green tissue on developing buds and repeat application during rains or immediately after until and including first cover. Later applications may be made as needed but no later than 1 day before harvest. Avoid application during periods of high temperature (90° F. or above) particularly when humidity is also high, so as to prevent possible fruit and foliage injury.

PEACHES - For control of brown rot (blossom blight) begin applications at time of balloon pink stage and continue with repeated applications at 4 day intervals until first cover. Later applications may be made as needed but no later than 7 days before harvest.

PLUMS, RED TART AND SWEET CHERRIES - For control of brown rot (blossom blight) begin applications in pre-blossom stage and continue with repeated applications at 4 day intervals until petal fall. Do not apply 3 days before harvest.