

E-Z-FLO

MANEB-SEVIN* 6-3 DUST

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Maneb (Manganese ethylenebisdithiocarbamate)	6.0%
Carbaryl (1-Naphthyl-N-methylcarbamate)	3.0%

INERT INGREDIENTS:

	91.0%
TOTAL	100.0%

*Sevin Reg. TM Union Carbide Chemical Corporation.
U. S. Pat. Nos. 2,903,478; 3,009,855.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

CAUTION

Harmful if inhaled or swallowed. May cause irritation of eyes, nose, throat and skin. Do not take internally, avoid breathing dust. Avoid contact with the skin, eyes and clothing. In case of contact with the skin, wash thoroughly with water; for eyes, get medical attention. Change contaminated clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of foodstuffs, feed, feeding troughs and water receptacles. Keep away from fire or sparks. Never allow dust to become wet during storage, as it will cause a chemical change which reduces its effectiveness and creates flammable vapors.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Net Contents and Manufacturer
On Container

DIRECTIONS FOR USE

CARROTS: Apply 50 lbs. per acre for the control of: ^{6/15/57} early & late blight. Apply when plants are 6 weeks old or when disease threatens. Repeat every 7 to 10 days. Flea beetles, harlequin bug, leafhoppers, six spotted leafhopper - Apply when insects appear at 5-7 day intervals.

CUCUMBER: Apply at the rate of 33 pounds per acre for the control of: anthracnose and downy mildew - Apply when plants begin to run or when disease threatens. Repeat at 5-7 day intervals as needed. Under severe conditions shorten intervals to 3 to 5 days. Cucumber beetle, flea beetle, leafhoppers, squash bug, pickleworm and melon-worm - Apply when insects or insect damage appears, repeat at 7 to 10 day interval. Avoid excessive application. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not apply within 5 days of harvest.

LETTUCE: Apply at the rate of 50 pounds per acre: downy mildew - Apply when disease appears, repeat at 7 to 10 day intervals under average conditions, 3 to 5 days when disease conditions are severe. Do not apply within 10 days of harvest and remove residues from head lettuce by stripping and trimming. Do not apply within 14 days to leaf lettuce and remove residues by washing or other effective means.

SUGAR BEETS: For control of Cercospora leaf spot, leafhoppers, armyworms, webworms and grasshoppers. Apply when conditions warrant and repeat at 7 to 10 day intervals, as needed. Apply at the rate of 40 pounds per acre. Do not apply within 14 days of harvest. Do not feed treated tops to livestock.

TOMATOES: Apply at the rate of 50 pounds per acre for control of early and late blight, Septoria leaf spot, gray leaf spot, anthracnose, Colorado potato beetle, leafhoppers, flea beetles, tomato fruit worm, fall armyworm, tomato horn worm, European corn borer, stink bugs, lace bugs and Lygus bugs. Begin applications when fruit of first cluster is well formed and repeat at 7 to 10 day intervals as needed. (Do not use on young seedlings in greenhouses as injury may occur.) Do not apply within 5 days of harvest.