

# E-Z FLO ALFALFA WEEVIL LARVAE SPRAY



Net Contents 5 Gallons  
(Contains 2 Lbs. Malathion and 3 Lbs. Methoxychlor Per Gal.)

## CAUTION: KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANELS FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS	
Malathion*	23.40%
Methoxychlor**, technical	23.40%
Xylene	46.97%
INERT INGREDIENTS	6.23%
	100.00%

The manufacturer makes no warranties, express or implied, concerning this product or its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

12-5


EPA Est. 635-MI-1; OH-1

E.P.A. Reg. No. 635-364

SOLD BY

## E-Z FLO CHEMICAL COMPANY

Division of Kirsto Company  
Lansing, Michigan • Columbus, Ohio



## DIRECTIONS FOR USE

For Use on Alfalfa- For the control of alfalfa weevil larvae, grasshopper, aphids, including spotted alfalfa aphid, lygus bugs, leafhoppers, spider mites, spittlebugs, vetch bruchids on alfalfa, apply 2 quarts of E-Z-Flo Alfalfa Weevil Control per acre in sufficient water for good coverage as a diluted spray.

For the control of Armyworms, apply 5 pints per acre.

Repeat applications as necessary. Apply this product to alfalfa in bloom only in the evening or early morning when bees are not working in the field or are not hanging on the outside of the hives.

Do not cut or graze within 7 days of application.

\*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

\*\*Equivalent to 20.59% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 2.81% other isomers and related compounds.

## CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with the skin. Do not breathe spray. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wash thoroughly with soap and water after use. Avoid contamination of feed and food-stuffs. Keep away from children and domestic animals.

NOTE TO PHYSICIAN: Give atropine preferably by injection, grains 1/30 to 1/15 (2 to 4 mg.) at once and every hour or oftener as needed to achieve and maintain full atropinization.

## USE PRECAUTIONS

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Highly toxic to fish. Do not contaminate any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use, pour, spill or store near heat or open flame.

Destroy Container When Empty.  
Never Re-use.