




# E-Z FLO TOBACCO FORMULA 1-3



**ACTIVE INGREDIENTS:**

<b>Parathion*</b>	<b>1.0%</b>
<b>Endosulfan**</b>	<b>3.0%</b>
<b>INERT INGREDIENTS:</b>	<b>96.0%</b>
	<b>100.0%</b>

\*O,O-Dimethyl diisopropyl phosphorothioate  
 \*\*Hexachlorocyclohexane 2, 4, 3-triazolodioxathiepin oxide. Thioran® a registered trademark of Canadian Hoechst Limited. U.S. Patent No. 2,799,685

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

This Tobacco Formula 1-3 is formulated on a non alkaline, sterilized tobacco base carrier. It is compounded especially for use on tobacco for the control of Aphids, Budworms, Cabbage Loopers, Hornworms, Cutworms, Vegetable Weevils and certain other insects.

## DIRECTIONS FOR USE

IN PLANT BED	NOTE	When and Where to Apply	Safety Restrictions
<b>Aphids and weevils</b>	Apply at 7 day intervals following transplanting and continue to end of growing season. Do not dust so as to obtain maximum coverage of the plant and plant parts. Use for aphid and budworm control.	Apply at 7 day intervals following transplanting and continue to end of growing season. Do not dust so as to obtain maximum coverage of the plant and plant parts. Use for aphid and budworm control.	Do not apply parathion within 5 days of transplanting.
<b>Cutworms</b>	Apply directly following transplanting.	Apply directly following transplanting.	
<b>Cabbage Looper and pinbugs</b>	Apply at 7 day intervals following transplanting and continue to end of growing season. Do not dust so as to obtain maximum coverage of the plant and plant parts. Use for aphid and budworm control.	Apply at 7 day intervals following transplanting and continue to end of growing season. Do not dust so as to obtain maximum coverage of the plant and plant parts. Use for aphid and budworm control.	

IN FIELD OR SHADE	NOTE
<b>Aphids</b>	Apply at 7 day intervals following transplanting and continue to end of growing season. Do not dust so as to obtain maximum coverage of the plant and plant parts. Use for aphid and budworm control.
<b>Budworms, Turnworms, and cabbage loopers</b>	Apply at 7 day intervals following transplanting and continue to end of growing season. Do not dust so as to obtain maximum coverage of the plant and plant parts. Use for aphid and budworm control.
<b>Cutworms</b>	Apply directly following transplanting.

**CAUTION:** No application should be made within 5 days of pruning tobacco or 15 days of cutting. Avoid plant parts coming in contact with the skin of those persons engaged in cutting or pruning the crop.