

E-Z FLO PHOSDRIN* 4 EC

(Contains 4 Lbs. of Active Ingredients Per Gallon)


DANGER

KEEP OUT OF THE REACH OF CHILDREN

See Side Panel for Additional Warnings and Antidote Statements



POISON



ACTIVE INGREDIENTS:		
Alpha isomer,2-carbomethoxy-1-methylvinyl dimethyl phosphate**	27.84%	
Related Compounds*	18.56%	
Xylene	41.57%	
INERT INGREDIENTS:	12.03%	
	100.00%	

**Equivalent to 46.4% Phosarin Insecticide.
*Shell Reg. Trademark.
Do Not Store Below 0° F.

Seller guarantees and warrants that the above stated information is true and correct and that the product is as described and labeled. Seller does not warrant that the product will be effective against all insects, diseases, or weeds. Seller's liability is limited to the purchase price of the product. Seller does not warrant that the product will be effective against all insects, diseases, or weeds. Seller's liability is limited to the purchase price of the product.

DIRECTIONS FOR USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

CONTAINER DISPOSAL: BEFORE DISCARDING THIS CONTAINER, FILL IT WITH WATER, ADDING 3 TABLETS OF A CAUSTIC (ALKALINE) SUBSTANCE, SUCH AS WASHING SODA, SODIUM CARBONATE, OR LIME. FILL WITH THE CONTAINER WITH WATER AND CAUSTIC TO STAND OVERNIGHT. BURY THE WASHINGS. EMPTY DRUMS MUST NOT BE USED IN ANY CONNECTION WITH FEED, FOOD OR DRINKING WATER. FOR 5 GALLON SIZE WASH, PERFORATE AND BURY CONTAINER WHEN EMPTY. NEVER RE-USE.

Do not apply later than 1 day before harvest. Do not spray on or near food. Apply in form of spray. Repeat applications when insects first appear. Repeat applications. Do not spray on or near food. Apply in form of spray. Repeat applications when insects first appear. Repeat applications. Do not spray on or near food. Apply in form of spray. Repeat applications when insects first appear. Repeat applications.

ALFALFA, CLOVER: Aphids, Alfalfa Caterpillar, Alfalfa Weevil, Lygus Bugs, Red Spider Mites, Grasshoppers, Leafhoppers, Cutworms (Climbing), Red Spider Mites, Lygus Bugs, Alfalfa Weevil, Lygus Bugs, Red Spider Mites, Grasshoppers, Leafhoppers, Cutworms (Climbing).

APPLES, PEACHES, PEAR, PLUMS, CHERRIES: Aphids, Red Spider Mites, Lygus Bugs, Red Banded Leaf Roller, 16 to 24 pt per 100 gals. water. Do not use with Chlorpyrifos. Do not apply more than 600 gals. per acre per season. Do not apply more than 600 gals. per acre per application on apples and pears.

BEANS, PEAS, POTATOES, CUCUMBERS, TOMATOES: Aphids, Red Spider Mites, Lygus Bugs, Cutworms (Climbing), Grasshoppers, Leafhoppers, Red Spider Mites, Lygus Bugs, Cutworms (Climbing), Grasshoppers, Leafhoppers, Red Spider Mites. **CAUTION** Do not exceed 16 pt per

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, LETTUCE, SPINACH, TURNIPS, BEETS, CARROTS, CELERY, EGGPLANT, PEPPERS: Aphids, Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt Marsh Caterpillar, Red Spider Mites, Cutworm (Climbing), Dipterous Leafminer (Adults), Lygus Bugs, Corn Earworm, False Chinch Bug. Do not apply to lettuce within 2 days of harvest. Do not apply to carrots, celery, eggplant, peppers, turnips, beets, and to spinach within 1 day of harvest.

CANTALOUPE, HONEYDEW MELON, MUSKMELONS, WATERMELONS, SUMMER SQUASH: Aphids, Cabbage Looper, Dipterous Leafminer (Adult), Leafhoppers, Lygus Bugs, Red Spider Mites, False Chinch Bug, Salt Marsh Caterpillars, Cutworms (Climbing), Grasshoppers. Do not apply to watermelons within 1 day of harvest.

GRAPES: Aphids, 16 to 24 pt in water to cover 1 acre. Leaf Folder, Leafminer, Red Spider Mites, Red Banded Leaf

APPLES, PEACHES, PEARS, PLUMS, CHERRIES: Aphids, Red Spider Mites — 1/4 to 1/2 pint per 100 gals. water. Grasshoppers, Lygus Bugs, Red Banded Leaf Roller — 1/2 to 3/4 pint per 100 gals. water. Do not apply to Cherries within 2 days of harvest. Do not apply more than 600 gals. per acre per application on cherries, peaches, and plums, nor more than 800 gals. per acre per application on apples and pears.

BEANS, PEAS, POTATOES, CUCUMBERS, TOMATOES: Aphids — 1/4 to 1/2 pint in water to cover 1 acre. Grasshoppers, Leafhoppers, Red Spider Mites, Mexican Bean Beetle (except potatoes) — 1/2 to 1 pint in water to cover 1 acre. **CAUTION:** Do not exceed 1/2 pint per acre.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, LETTUCE, SPINACH, TURNIPS, BEETS, CARROTS, CELERY, EGGPLANT, PEPPERS: Aphids — 1/4 to 1/2 pint in water to cover 1 acre. Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt Marsh Caterpillar, Red Spider Mites, Cutworm (Climbing), Dipterous Leafminer (Adults), Lygus Bugs, Corn Earworm, False Chinch Bug — 1/2 to 1 pint in water to cover 1 acre. Do not apply to Lettuce within 2 days of harvest, to Brussels Sprouts, Cauliflower, Celery, Collards, Beets, Kale and Turnips within 3 days and to Spinach within 4 days of harvest.

CANTALOUPES, HONEYDEW MELON, MUSKMELONS, WATERMELONS, SUMMER SQUASH: Aphids — 1/4 to 1/2 pint in water to cover 1 acre. Cabbage Looper, Dipterous Leafminer (Adult), Leafhoppers, Lygus Bugs, Red Spider Mites, False Chinch Bug, Salt Marsh Caterpillars, Cutworms (Climbing), Grasshoppers — 1/2 to 1 pint in water to cover 1 acre.

GRAPES: Aphids — 1/2 to 1 pint in water to cover 1 acre. Leaf Folder, Leafhopper, Red Spider Mites, Red Banded Leaf Roller, Lygus Bugs — 1 to 2 pints per acre. Do not apply within 2 days of harvest.

ONIONS (Including Green Onions): Thrips, Cutworms (Climbing) — 1/2 to 1 pint in water to cover 1 acre.

RASPBERRIES: Aphids — 1/4 to 1/2 pint per 100 gals. water. Red Spider Mites, Leafhoppers, Fruit Tree Leafroller — 1/2 to 3/4 pint per 100 gals. water. Do not treat within 3 days of harvest.

02-23

53-263-880

U.S.D.A. Reg. No. 635 376

DIRECTIONS FOR USE (Continued)

STRAWBERRIES: Aphids, Red Spider Mites, Thrips — 1/4 to 1/2 pint per 100 gals. water (200 gals. diluted spray per acre). Grasshoppers, Strawberry Leaf Roller, Salt Marsh Caterpillar, Lygus Bugs — 1/2 to 1 pint per 100 gals. water per acre.

USE PRECAUTIONS

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Toxic to fish and wildlife. Do not contaminate any body of water or apply to any area not specified in the label. Birds feeding on treated areas may be killed. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

WARNING

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

APPLES, PEACHES, PEARs, PLUMs, CHERRIES: Aphids, Red Spider Mites. 16 to 24 pint per 100 gal. water. Grasshoppers, Lygus Bugs, Red-Banded Leaf Roller. 16 to 24 pint per 100 gal. water. Do not apply to Cherries within 2 days of harvest. Do not apply more than 600 gal. per acre per application on apples and pears.

BEANS, PEAS, POTATOES, CUCUMBERS, TOMATOES: Aphids, Red Spider Mites, Mites, Beetles. CAUTION. Do not exceed 1/2 pint per acre.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, LETTUCE, SPINACH, TURNIPS, BEETS, CARROTS, CELERY, EGGPLANT, PEPPERS: Aphids, Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt Marsh Caterpillar, Red Spider Mites, Cutworm, Tomlinson, Cabbage Leafminer (Adults), Lygus Bugs, Corn Earworm, Colorado Potato Beetle, Flea Beetles, Wireworms, and the Spiny Nematode.

CANTALOUPEs, HONEYDEW MELON, MUSKMELONS, WATERMELONS, SUMMER SQUASH: Aphids, Cabbage Looper, Imported Cabbage Worm, Grasshoppers, Leafhoppers, Salt Marsh Caterpillar, Red Spider Mites, Cutworm, Tomlinson, Cabbage Leafminer, Bug, Salt Marsh Waterpillars, Cutworms (Tomlinson), Colorado Potato Beetle.

GRAPES: Aphids, Red Spider Mites, Mites, Beetles. Do not apply to grapes within 2 weeks of harvest.

ONIONS (Including Green Onions): Aphids, Red Spider Mites, Mites, Beetles.

RASPBERRIES: Aphids, Red Spider Mites, Mites, Beetles. Do not apply to raspberries within 2 weeks of harvest.

02 23 53-263-880

DIRECTIONS FOR USE (Continued)

STRAWBERRIES: Aphids, Red Spider Mites, Mites, Beetles. Do not apply to strawberries within 2 weeks of harvest.

USE PRECAUTIONS

Do not apply under conditions of extreme heat or drought. Do not apply to plants that are stressed or damaged. Do not apply to plants that are in bloom. Do not apply to plants that are in fruit. Do not apply to plants that are in seed. Do not apply to plants that are in storage. Do not apply to plants that are in transit. Do not apply to plants that are in display. Do not apply to plants that are in use.

DESTROY CONTAINER WHEN EMPTY NEVER REUSE