

# E-Z FLO BHC 12-W

**CAUTION:**  
KEEP OUT OF REACH OF CHILDREN

**ACTIVE INGREDIENTS:**

Gamma Isomer of Benzene Hexachloride (BHC).....12.0%  
Other Isomers of Benzene Hexachloride (BHC).....36.0%

**INERT INGREDIENTS:**.....52.0%  
Total 100.0%

**CAUTION**

Harmful if Swallowed. May be absorbed through skin. May produce irritation of skin and eyes. Wash thoroughly after handling. Avoid contact with skin and eyes. Wear respirator and goggles. Do not breathe dust or spray mist. Avoid contamination of food, feed, seed, and fertilizer.

52-28      EPA Est. 635-MI-1; OH-1      E.P.A. Reg.No. 635-378

**DIRECTIONS FOR USE**

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank, or pre-mix in a bucket half filled with water and pour into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

ACCEPTED  
AUG 26 1975  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 635-378

**PEAR & APPLE:** Aphids - Use 1/2 to 3 lbs. in 100 gals. water. May be used as late as petal fall at the 2 lb. dosage for the control of Rosy Apple Aphids. Do not apply more than 12 lbs. per acre application. Do not apply within 60 days of harvest, or after the fruit starts to form.

**PEACH:** Plum Curculio - Use 1/2 to 2 lbs. in 100 gals. water. Do not apply more than 8 lbs. per acre application. Make first application when half of shucks have fallen, second application when all of shucks have fallen, and third application about 7 days later. Do not use within 60 days of harvest on peaches.

**STRAWBERRIES:** Spittle Bugs, Strawberry Weevils - Use 1/2 to 1 lb. in 100 gals. of water. Do not apply more than 16 lbs. per acre application. Do not apply after the edible parts start to form.

**USE PRECAUTIONS**

Do not apply on feed or forage crops to be fed to dairy animals or to animals being finished for slaughter.

Do not apply to crops in rotation with root crops, or use in soils that are to be planted with these crops since off-flavor will result.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Consult your State Agricultural Experiment Station for additional information, as the timing may vary with local conditions.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not apply during bloom when honey bees or other pollinators are active. Avoid drift to adjoining food and forage crops or crops in bloom.

This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Not Contained