

Net Content 50 lbs. per container
On Container

ACCEPTED
apr 8 1971
635-388
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO. 635-388 SUBJECT
D ATTACHED COMMENTS.

E-Z-FLO

Parathion, Maneb, Copper,

1.5-3.8-4.5 DUST

WARNING: KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

| | |
|---|------|
| *Copper (expressed as metallic copper) | 4.5% |
| Maneb (Manganese ethylenebisdithiocarbamate) | 3.8% |
| Parathion (0,0-Diethyl 0-p-Nitrophenyl Thiophosphate) | 1.5% |

INERT INGREDIENTS:

90.2%
TOTAL 100.0%

*From Basic Copper Sulphate

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

CELERY: Early blight (Cercospora), Late blight (Septoria), Celery worms and aphids. Apply when plants are established in the field; repeat as frequently as required (every 7 days under average conditions or every 3-5 days under severe insect or disease conditions). Remove excess residues by stripping, trimming and washing. Do not apply within 21 days of harvest. Apply 33 lbs. per acre.

ONIONS: Downy mildew, purple blotch (blight), Blast (Botrytis leaf blight) and thrips. Apply when first disease or thrips appear. Repeat weekly as needed. Do not apply within 15 days of harvest. Apply 30-35 lbs. per acre.

POTATOES: Early blight (Alternaria), Late blight (Phytophthora), Colorado potato beetle, Flea beetle, Leaf hopper, Serpentine leaf miner, Grasshoppers, Whitefly and aphids. Apply as soon as plants are 2-6 inches high; repeat at 5-10 day intervals as long as blights or insects threaten. Under severe disease conditions, use the shorter interval or the higher dosage. Apply 30-35 lbs. per acre. Do not apply within 5 days of harvest.

WARNING

May be fatal if swallowed, inhaled or absorbed through the skin. Rapidly absorbed through the skin. Do not get on the skin, in the eyes or on clothing. Do not breathe dust. Wear a respirator which has been tested by the U.S. Department of Agriculture for parathion protection. Wear waterproof or protective clothing, natural rubber gloves and hat. Keep away from all feed and food products. Wash hands, arms and face with soap and water thoroughly after application or before eating or smoking. Wash contaminated clothing with hot water and soap before reusing.

PHYSICIAN'S NOTE: Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine, preferably by injection, grains 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. 2-PAM is also antidotal and may be administered in conjunction with atropine. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

Not for use or storage in or around the home.

Keep all unprotected persons or animals out of the operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residue has dissipated. Do not dust plants during pollinations and/or flowering periods.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Birds feeding on treated areas may be killed.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dispose of wastes and spillage by burying in non-crop land away from water supplies, containers should be disposed of by burning (stay out of smoke).