

# E-Z FLO TOBACCO FORMULA 1-3-10

### (CONTAINS 1% PARATHION, 3% ENDOSULFAN AND 10% DITHANE<sup>(11)</sup>)

## **DANGER:** KEEP OUT OF THE REACH OF CHILDREN



ACTIVE MICHELIA	
Parathion (O, O-diethyl O-p-nitrophenyl	
thiophosphate)	1.0%
Endosulfan* (Hexachlorohexahydromethano-2,	
4, 3-benzodioxathiepin oxide)	3.0%
Zineb (Zinc ethylene bisdithiocarbamate) * *	6.5%
INERT INGREDIENTS:	89.5%
*Thiodan®	100.0%

\*Thiodan(R)

\*\*Equivalent to 10% Dithane.

T) Reg. Trademark of Rohm and Haas Company.

(Reg. Trademark of Canadian Hoechst Limited, U.S. Patent No. 2,799,685.

### DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

This "Tobacco Formula" Dust is formulated on a non-alkaline, sterilized tobacco base carrier It is compounded especially for use on tobacco for the control of Aphids, Budworms Cabbage Loopers, Hornworms, Cutworms, Vegetable Weevils, and certain other insects, as well as for the control of Tobacco Blue Mold. "Do not apply within 5 days of priming or 15 days of cutting. Avoid having plant juices coming into contact with the skin."

Blue Mold: In the Plant Bed: Make first application when plants are about the size of a dime using 10 to 1 lbs. per 100 sq. yards and increase to 31/2 lbs. as plants mature, Following applications twice per week until plants free of blue mold are assured for transplanting. In the Field: Apply on newly-set plants at the rate of 7-8 lbs, per acre, increasing to 18 to 20 lbs. as the plants mature. Repeat applications at 14-day intervals or after each rain. Do not apply within 5 days of priming or 15 days of cutting.

Dosage per acre or as otherwise Insect indicated		When and where to apply	Safety Restrictions	
IN PLANT BED Aphids and vegetable weevils		On foliage at seven-day intervals be- ginning three weeks prior to trans- planting for aphids or as needed for vegetable weevils	Do not apply Parathion within 5 days before transplanting.  Should be applied only by a trained operator.	

	osage per acre r as otherwise indicated	When and where to apply	Safety Restrictions
Cutworms	0.4-0.7 lb./ 100 sq. yd.	On foliage as needed.	
Garden Slug and pillbugs	s. 0.4-0.7 lb./ 100 sq. yd.	These pests hide under boards, wall flaps and under debris in the plant hed. Applications should be directed at hiding places at needed.	
IN FIELD OR SHADE Aphids	20·25-lb.	On foliage at seven-day intervals beginning immediately following transplanting and continuing to end of priming season. Direct dust so as to obtain maximum coverage of the plant and particularly the bud for aphid and budworm control.	Should be applied only by a trained operator
Budworms, Hornworms, and cabbage loopers		On foliage at seven-day intervals be- ginning immediately following trans- planting and continuing to end of priming season. Direct dust so as to obtain maximum coverage of the plant and particularly the bud for aphid and budworm control.	Should be applied only by a trained operator
Cutworms	18-21 lb.	Apply directly following transplanting.	

#### **WARNING:**

The Statement, they bus to see the both there is at size of intrough the law of Mean and a corner cours of the property of the period of the specific region of the and the second section of the second second second second second section is a second s the control of the category of the category of the control of the category of Control of the contro

• •

Cantinner Disposa THE AST ASD TRUNCHS

CALL A PHYSICIAN AT MET

en de la companya de la co

02-2

# E-Z FLO TOBACCO FORMULA 1-3-10

## (CONTAINS % PARATHION, 3% ENDOSULFAN AND 10% DITHANE")

	) 2 4	A LANGE			
	!	1. 1.	٠,		•
		y v			
ACTIVE INGREDIENTS.					
Parathion (O, O-diet	hyl	O-p-nitro	phenyl		
thiophosphate)					1.0°°
Endosulfan* (Hexac	hlor	ohexahy	drometh	ano-2,	ı
4, 3-benzodioxa	athie	pin oxid	le)		3.0°°
Zineb (Zinc ethylene bisdithiocarbamate) **			6.5°o		
INERT INGREDIENTS:					89.5°
					100.0%
				·	

#### DESTROY CONTAINER WHEN EMPTY, NEVER RE-USE

#### DIRECTIONS FOR USE

N -11 A - 1 - 2 - 3 ft.

	sali je kire. Postorova	After our work	5,00
	10.1		here is no
	C4011	rots agents bested	
	) ; P 7 %	Trace pasts to deconder to come was tops and under debres ender part of App of ms stoud to discould be discould be also as	
IN FIELD OR SHADE	: : .	the specific environments, letters entry the second steel of the second steel of the second s	in until be appeared as a
		The second of th	
· · · · · · · · · · · · · · · · · · ·	٠,	A second of the second of t	
231		And the second	

1, .