E-Z-Flo Thiodan[®] Copper 1.5 - 6 Dust

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

93.1%

ACTIVE INGREDIENTS:

Endosulfan (hexachlorohexahydromethano-2,4, 3benzodioxathiepin oxide) 1.5% Copper-Zinc-Chromate Complex (15 CuO-10ZnO-6CrO₃-25 H₂O) 5.4% Copper expressed as metallic 1.68%


Copper expressed as metallic 1.68%
Zinc expressed as metallic 1.16%
Chromium expressed as metallic 0.55%
INERT INGREDIENT:

Total 100.0%

*Registered trademark of Canadian Hoechst Limited. U. S. Patent No. 2,799,685.

*Equivalent to 6% Miller 658.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.



POTATOES—Colorado potato beetle, Potato flea beetle, Potato leafhopper, Aphids, Southern Armyworm, Green stink bug. Leaf-footed plant bug and Potato tuberworm, Early and Late Blight. Apply 60-65 pounds per acre when insects first appear or when potatoes are 4-6 inches high (whichever is first) and continue on a 7 to 10 day schedule. Shorten the interval if serious disease or insect conditions prevail. Workers entering treated areas the day of application should wear protective clothing. No time limitations. Do not plant root crops other than carrots, potatoes, and sweet potatoes as follow-up crops.

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust mist. Avoid contact with skin, eyes or clothing. Avoid contamination of feed or foodstuffs. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing.

Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Birds and other wildlife feeding on the treated area may be harmed.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

ontents and Manufacturar On Container