

# E-Z FLO LINDANE 25 W

**WARNING:**  
**KEEP OUT OF THE REACH OF CHILDREN**

REC'D  
 Oct 4 1973  
 U.S. DEPARTMENT OF AGRICULTURE  
 REGISTERED

**ACTIVE INGREDIENTS:**

**Gamma isomer of benzene hexachloride**  
 (from lindane) ..... **25.0%**

**INERT INGREDIENTS:** ..... **75.0%**  
 ..... **100.0%**

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

## DIRECTIONS FOR USE

**FARM PREMISES**—Exempt Poultry, Horses, and Goats—Mix 1 lb. per 100 gal. water. Before applying spray, remove livestock and feed from premises and feed and water troughs. Do not contaminate milk rooms, milking equipment, feed, or drinking water. Do not use on premises where children play.

**FOR FLIES AND MOSQUITOES**—Apply as a coarse spray at rate of 1 gal. per 500 sq. ft. or wet all surfaces to point of run-off. Also spray manure piles and other breeding places. Repeat as necessary.

**FOR ROACH, ANT, BEDBUG AND TICK CONTROL**—Make thorough application to corners, stalls, and other areas where pests are frequently found. Repeat as necessary.

**FARM ANIMALS**—Exempt Dairy Animals—Do not use on lactating dairy animals.

**SARCOPTIC MANGE MITES, LICE, TICKS**—Do not spray on animals within 30 days of slaughter. Do not dip animals within 10 days of slaughter. Do not use spray containing more than 0.06% gamma. Do not use on young animals including calves under 3 months of age. Other immature animals may be injured by treatment. Spray or dip application to young animals over 3 months of age should be limited to 0.03% concentration of Lindane.

**SARCOPTIC MANGE MITES ON BEEF CATTLE**—Use 1 lb. per 100 gal. water (0.03-0.045% concentration) on animals 10 months of age and over. Use lower dosage on animals under 6 months. Make thorough application.

**LICE AND TICKS ON BEEF CATTLE, SHEEP, HORSES AND GOATS**—Use 1 lb. per 100 gal. water (0.03-0.045% concentration) on animals 10 months of age and over. Use lower dosage on animals under 6 months. Repeat treatments at 7-10 day intervals may be necessary to prevent reinfestation.

**SARCOPTIC MANGE MITES AND LICE ON GOATS**—Use 2 lbs. per 100 gal. water (0.06% gamma concentration) on animals 10 months of age and over. Repeat treatments one week apart.

**CAUTION**—Do not treat swine with this product. Do not treat swine at least three weeks after. Avoid treating animals near or during milking. Avoid treating swine that are milk animals.

**INSECT CONTROL ON FLOWERS AND VEGETABLES:**

**CUCUMBERS, CANTALOUPE, SQUASH AND PUMPKINS**—For control of cucumber beetles, thrips use 1.2 to 2 lbs. per acre. Aphids, melonworms, leaf miner—use 1.6 to 2 lbs. per acre. Do not use more than 2 pounds per acre. Remove excess residues from cucumbers and squash. Do not use more than 2 pounds E-Z FLO Lindane W-25 per acre. Do not apply to cucumbers and squash within one day of harvest.

**SPINACH**—Aphids, leafminers—Use 1.6 to 3.0 lbs. per acre. Apply only in seedling stage.

**CELERY**—Aphids, thrips, tarnished plant bugs—use 1.2 to 1.6 lbs. per acre. Do not apply after the celery is half grown or after it begins to bunch.

**CABBAGE AND BROCCOLI**—Aphids, thrips—use 0.8 to 1.2 lbs. per acre. Do not apply after edible parts start to form. Do not use more than 3 pounds E-Z FLO Lindane W-25 per acre.

**GLADIOLUS**—Thrips—1 lb. per 100 gal. water.

**INSECT CONTROL ON TREES AND SHRUBS:**

**FLOWERING FRUITS, ELM, BEECH, MAPLE, SYCAMORE, BIRCH**—Aphids 1 lb. per 100 gal. water. Apply when new growth is about 2 inches long.

**AZALEA, RHODENDRON, OAK AND SYCAMORE**—Lace bugs—1 lb. per 100 gal. water. Apply just after hatching when small nymphs can be seen. Second spray 10 days later.

**PINES**—Pine bark aphid—1 lb. per 100 gal. water. Apply when immature aphids are present.

## WARNING

**MAY BE FATAL IF SWALLOWED.** May be Absorbed through skin. Do not breathe spray mist. Avoid contact with skin or eyes. In case of contact, wash immediately with soap and water. For eyes, flush with water and get medical attention. Avoid contamination of feed and food stuffs. Do not use on cats or other household pets nor on poultry or humans.

**LINDANE W-25** should not be applied under conditions which might result in a toxic residue, or residual odor or taste remaining on foodstuffs or edible portions of treated crops or forage at harvest time. Do not use on forage crops to be fed to dairy animals or animals being finished for slaughter. Do not use in poultry houses or dairy barns. Do not treat storage piles of grain to be sold immediately or shipped interstate. Do not use at higher concentrations than recommended on this label.

Toxic to fish and wildlife. Do not contaminate any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees. Do not use for treatment of bees for varroa mites. For more information may be obtained from your local Agricultural Extension Service.

Do not reuse empty container. Do not mix with other pesticides. Wash thoroughly. Stay away from smoke or fumes.

Net Contents and Manufacturer  
 On Container