# E-Z FLO MCPP

## CAUTION: KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:
Potassium salt of 2-(4-Chloro-2-methylphenexy)
propionic acid\*
INERT INGREDIENTS:

68.9%

\*Equivalent to 25.4% 2-(4-Chloro-2-methylphenoxy) propionic acid.

NOTICE OF WARRANTY: Seller makes ne warranty of any kind concerning the use of this product. User assumes all risk for use or handling whether in accordance with directions or not.

#### SOLD BY

#### E-Z FLO CHEMICAL COMPANY DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA COLUMBUS, OHIO LANSING, MICHIGAN LEXINGTON, KENTUCKY

72-13R

53-263-843

U.S.D.A. Reg. No. 635-403

100.0%

## **DIRECTIONS FOR USE**

MCPP Herbicide will not injure bent grasses, established lawns or turf grasses when used as directed. Controls chickweed, clover, plantain, black medical and knotweed in turf grasses. Apply when weeds are young and growing vigorously. Do not apply to bent grasses in the heat of the Summer. Do not mow for several days before or after treatment. Do not water for one day following treatment.

### ESTABLISHED GRASSES (Fairways and Lawns)

Apply 1½ to 2¼ fluid ounces per 1,000 square feet in 2½ gallons of water (4 to 6 pints in at least 25 gallons of water per acre). Do not spray during long dry spells.

**NOTE:** If dandelions are a problem add 2,4-D at the usual rate except on bent grasses where the rate of 2,4-D should not exceed 4 oz. acid equivalent per acre.

#### ESTABLISHED GREENS

Apply 1½ fluid ounces per 1,000 square feet in 2½ gallons of water (4 pints in at least 25 gallons of water per acre) for control of mouse ear chickweed, black medic, knotweed and plantains.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

JUL 2 6 1967

FUNDER THE MAL NAME TIGIDE FUNDICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REDISTERED UNDER NO. 6 10 3 SUBJECT TO ATTACHED COMMENTS.

ACCEPTED

## **CAUTION**

Keep out of reach of children. Avoid contact with skin, eyes and clothing. Wash hands after use. Keep in original container in storage.

Do not allow spray, spray mist, visible or invisible vapors, or air containing mist or vapors to contact plants, other than those being sprayed. If 2,4-D or MCPP susceptible plants are growing within a distance of ½ mile from the area being sprayed, make application only when there is a light breeze away from the susceptible plants. Spray drift can be reduced by using high volume sprays under low pressure, coarse droplet nozzles and boom nozzle drops. Susceptible crops include grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, cauliflower and cabbages, box elder trees, as well as many other field vegetables, flowers, ornamental and fruit crops.

Do not contaminate streams, lakes, ponds, irrigation water, or water used for livestock or domestic purposes.

Do not store open containers near seeds, feeds, fertilizers or pesticides including insecticides, fungicides and weed killers.

Sprayers which have contained MCPP should not be used to apply other pesticides to 2,4-D or MCPP susceptible crops.

Rinse equipment and containers and dispose of wastes and spillage by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning (keep out of smoke).

MARRANTY STATEMENT PRESENTLY UNDER STUDY AND WILL BE REVISED