

ACCEPTED
 May 18 1971 635-407
 UNDER THE FEDERAL INSECTICIDE ACT
 REGISTERED PRODUCT
 EPA REG. NO. 635-407

1-Z FLO HEPTACHLOR 20.3

WARNING:
KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENTS:

Heptachlor*	20.3%
Related Compounds	7.8%
INERT INGREDIENTS:	72.2%
	0.0%

*Heptachlorotetrahydro-4,7-methanoindene.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE
 DIRECTIONS FOR SOIL TREATMENT

BROADCAST: Make preplant broadcast application with a combination drill or fertilizer spreader, or by airplane, to cover the soil surface uniformly. Follow promptly with cultivation to mix the insecticide into the top 3 to 6 inches of soil.

RECOMMENDATIONS

BERRIES (Blackberries, boysenberries, blueberries, dewberries and raspberries): Strawberry root weevil larvae (Brachyrhinus) - apply 20 lbs per acre. Do not apply after edible parts start to form.
CORN: Corn rootworm-apply 4 to 5 lbs per acre. Cornfieldants, root aphids, seed corn maggots and seed corn beetles- apply 7.5 lbs. per acre. Cutworms - apply 5 to 10 lbs per acre. Asiatic garden beetle larvae, European chafer larvae, green June bug larvae, Japanese beetle grubs - apply 10 lbs. per acre.
PEPPERS and TOMATOES: Cutworms, seedcorn maggot, and wireworms - apply 10 lbs per acre. Do not apply after edible parts start to form.

SMALL GRAINS (Barley, Oats, Rye, and Wheat): White grubs and wireworms - apply 10 to 15 lbs per acre as a preplant soil application and disc into the top 3 to 6 inches of soil.
ORNAMENTAL TURF, LAWN, NURSERY AND GREENHOUSE SOILS:
 For use in turf (such as golf courses, athletic fields, large estate nurseries, cemeteries and park sod), and in nursery and greenhouse soils where ornamental plants are grown. Cover surface uniformly using a lawn fertilizer spreader. Treatment should be applied in the spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling. Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water, and grass has completely dried. For the control of Earwigs and Striped Grassworms, use 5 lbs. per acre; for the control of Ants and Cutworms, use 10 lbs. per acre; for the control of Amyworms, Asiatic Garden Beetle Grubs, European Chafer Grubs, Green June Beetle Grubs, Japanese Beetle Grubs, Lawn Chinch Bugs, Mole Crickets, Northern Masked Chafer Grubs, Pillbugs, Snailbugs, Sowbugs, and White Grubs, use 15 lbs. per acre.

WARNING

May be fatal if swallowed. Do not breathe vapor or dust. Do not get on skin. In case of contact wash immediately with soap and water. Do not contaminate feed and foodstuffs.
 Avoid dust drift onto adjoining food and forage crops or crops in bloom. Apply in early morning or evening when there is little or no wind. Do not use this product on feed or forage to be fed to dairy animals or to animals being finished for slaughter.
 To protect bees, do not apply when crops are in bloom.
 This product is poisonous to wildlife and fish. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on label. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.