

E-Z FLO 5% MALATHION DUST

CAUTION:

KEEP OUT OF THE REACH OF CHILDREN

ACTIVE INGREDIENT:

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) 5.0%

INERT INGREDIENTS: 95.0%
100.0%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

DIRECTIONS FOR USE

Applications should be made when insects first appear and continued at 7 to 10 day intervals until controlled. Do not dust plants during flowering and/or pollination periods.

ASPARAGUS- Thrips, Asparagus Beetles. Use 20-25 lbs. per acre. Do not apply within 1 day of harvest.

BEANS- Mexican Bean Beetles, Leafhoppers, Asparagus Beetles, Aphids, Cucumber Beetles. Use 30-35 lbs. per acre. Do not apply within 1 day of harvest.

CELERY- Aphids, Spider Mites. Use 20-30 lbs. per acre. Do not apply within 7 days of harvest.

CUCUMBERS, SQUASH, MELONS- Aphids, Spider Mites, Flea Beetles, Cucumber Beetles. Use 30-35 lbs. per acre. Do not apply to cucurbits unless plants are dry. Do not apply within 1 day of harvest.

CALIF. CABBAGE, CAULIFLOWER- Aphids, Imported Cabbageworms, Cabbage Looper. Use 20-25 lbs. per acre. Do not apply within 7 days of harvest.

ETTUCE- Aphids, Mites, Cabbage Looper, Leafhoppers. Use 20-25 lbs. per acre. Do not apply to lettuce within 7 days of harvest.

LEAFY VEGETABLES- Aphids, Spider Mites. Use 20-25 lbs. per acre. Do not apply within 7 days of harvest.

PEAS- Aphids, Spider Mites, Flea Beetles, Cucumber Beetles. Use 30-35 lbs. per acre. Do not apply within 1 day of harvest.

DIRECTIONS FOR USE

STRAWBERRIES- Lygus Bugs, Scabbiebugs, Aphids, Spider Mites, Flea Beetles, Leafhoppers, Strawberry Leafhoppers, White Flies. Use 20-40 lbs. per acre. Do not apply within 3 days of harvest.

ORNAMENTALS- Aphids, Mites, Plant Bugs, Scale, Thrips, Mealy Bugs, White Fly. Use 20-40 lbs. per acre. No days between last application and harvest.

BRUSSELS SPROUTS- Aphids. Use 30 lbs. per acre. Do not apply within 7 days of harvest.

CARROTS, RADISHES, PARSNIPS, SALSIFY, HORSE RADISH- Aphids. Use 30 lbs. per acre. Do not apply within 7 days of harvest.

ENDIVE- Aphids, Mites. Use 30 lbs. per acre. Do not apply within 7 days of harvest.

PUMPKINS- Aphids, Mites, Leafhoppers. Use 25-30 lbs. per acre. Do not apply within 3 days of harvest.

COTTON- Boll Weevil, Desert Spider Mite, Cotton Aphid, Leafhopper, White Flies, Thrips, Brown Cotton Leafworm, Cotton Leaf Perforator, Cotton Leaf Worm, Lygus Bug. Use 20-25 lbs. per acre. No time limitation.

TOBACCO- Aphids. Use 20-30 lbs. per acre. No time limitation.

CATTLE- Beef cattle and non-lactating dairy cattle. Hornflies apply on back and neck of the animals and repeat at 10-14 day intervals. For Lice control treat the animal thoroughly. Repeat in 10 days and thereafter as needed. For ticks, treat the entire animal thoroughly and repeat at 2 week intervals as needed.

Do not treat animals under one month of age. Do not apply to lactating dairy animals or within 2 weeks of freshening. Do not contaminate feed, food containers, and watering troughs.

CATTLE- Lactating Dairy Cattle. Horn Fly, apply 3 tablespoons of 5% Malathion dust per animal. Apply on back and neck of the animals and repeat applications at 10-14 day intervals. Do not apply to Dairy Cattle later than 5 hours before milking or during milking. Do not contaminate water troughs.

POULTRY- (Chickens, Ducks, Geese and Turkeys). To control Poultry Lice, Northern Fowl Mite, and Chicken Red Mite dust individual birds by means of a shaker can or hand duster. Repeat in 4 to 8 weeks when necessary. Premise Treatment. Apply 1 pound per 50 square feet of litter and floor space and to nests, roost and adjacent areas.

PIGS- For lice control, treat animal thoroughly and repeat in 10 days and thereafter as needed. Avoid contamination of feed, food containers, and watering troughs.

CAUTION

Avoid contact with eyes. Wash thoroughly with soap and water immediately if dust gets into eyes. If dust gets into mouth, spit out and wash mouth with water. Do not swallow. Do not breathe dust.

Keep away from children and pets. Do not use in or near living quarters.

Keep away from food and feed. Do not use in or near living quarters.

Do not use in or near living quarters. Do not use in or near living quarters.

DIRECTIONS FOR USE

Applications should be made when insects first appear and continued at 7 to 10 day intervals until controlled. Do not dust plants during flowering and/or pollination periods.

ASPARAGUS- Thrips, Asparagus Beetles: Use 20-25 lbs. per acre. Do not apply within 1 day of harvest.

BEANS- Mexican Bean Beetles, Leafhoppers, Japanese Beetles, Aphids, Cucumber Beetles: Use 30-35 lbs. per acre. Do not apply within 1 day of harvest.

CELERY- Aphids, Spider Mites: Use 25-30 lbs. per acre. Do not apply within 7 days of harvest.

CUCUMBERS, SQUASH, MELONS- Aphids, Spider Mites, Pickle Worms, Cucumber Beetle: Use 30-35 lbs. per acre. Do not apply to cucurbits unless plants are dry. Do not apply within 1 day of harvest.

KALE, CABBAGE, CAULIFLOWER- Aphids, Imported Cabbage Worms, Cabbage Looper: Use 30-35 lbs. per acre. Do not apply within 7 days of harvest.

LETTUCE- Aphids, Mites, Cabbage Loopers, Leafhopper: Use 25-30 lbs. per acre. Do not apply to head lettuce within 7 days of harvest, nor to leaf lettuce within 14 days of harvest.

PEAS- Pea Aphids, Pea Weevil: Use 25 lbs. per acre. Do not apply within 3 days of harvest.

PEAS (vines)- Pea Aphids, Pea Weevil: Use 25 lbs. per acre. Do not apply within 3 days of harvest nor within 7 days of harvest if forage is to be fed to livestock.

POTATOES- Aphids, Leafhoppers: Use 25-30 lbs. per acre. No time limitations.

SPINACH- Aphids: Use 30 lbs. per acre. Do not apply within 7 days of harvest.

TOMATOES- Tomato Russet Mite, Spider Mite, Aphids, Drosophila: Use 35-40 lbs. per acre. Do not apply within 1 day of harvest.

BLUEBERRIES- Blueberry Maggot, Japanese Beetles, Plum Curculio, Sharpnose Leafhopper: Use 30-40 lbs. per acre. Do not apply within 1 day of harvest.

RASPBERRIES- Aphids, Rose Scale, Mite, Thrips, Leafhoppers: Use 35 lbs. per acre. Do not apply within 1 day of harvest.


CATTLE- Beef ca
day intervals. Fo
entire animal thor

Do not treat anim
not contaminate fi

CATTLE- Lactati
of the animals and
or during milking.

POULTRY- (Chic
dust individual bi
Apply 1 pound pe

HOGS- For lice
feed, food contai



Avoid contact wit
handling this prod

6-35-4 NOTE TO PHYSIC

Keep away from c

Highly toxic to fi
when weather con
wastes. Apply th

This product is hig
from your Cooper

Avoid applicatio

411-22

CHILDREN

DIRECTIONS FOR USE

STRAWBERRIES- Lygus Bugs, Spittlebugs, Aphids, Spider Mite, Potato Leafhopper, Strawberry Leafroller, White Flies: Use 20-40 lbs. per acre. Do not apply within 3 days of harvest.

ORNAMENTALS- Aphids, Mites, Plant Bugs, Scale, Thrips, Mealy Bugs, White Fly: Use 20-40 lbs. per acre. No days between last application and harvest.

BRUSSELS SPROUTS- Aphids: Use 30 lbs. per acre. Do not apply within 7 days of harvest.

CARROTS, RADISHES, PARSNIPS, SALSIFY, HORSERADISH- Aphids: Use 30 lbs. per acre. Do not apply within 7 days of harvest.

ENDIVE- Aphids, Mites: Use 30 lbs. per acre. Do not apply within 7 days of harvest.

PUMPKINS- Aphids, Mites, Leafhoppers: Use 25-30 lbs. per acre. Do not apply within 3 days of harvest.

COTTON- Boll Weevil, Desert Spider Mite, Cotton Aphid, Leafhopper, White Flies, Thrips, Brown Cotton Leafworm, Cotton Leaf Perforator, Cotton Leaf Worm, Lygus Bug: Use 20-25 lbs. per acre. No time limitation.

TOBACCO- Aphids: Use 20-30 lbs. per acre. No time limitation.

CATTLE- Beef cattle and non-lactating dairy cattle: Hornflies, apply on back and neck of the animals and repeat at 10-14 day intervals. For Lice control treat the animal thoroughly. Repeat in 10 days and thereafter as needed. For ticks, treat the entire animal thoroughly and repeat at 2 week intervals as needed.

Do not treat animals under one month of age. Do not apply to lactating dairy animals or within 2 weeks of freshening. Do not contaminate feed, food containers, and watering troughs.

CATTLE- Lactating Dairy Cattle: Horn Fly, apply 3 tablespoons of 5% Malathion dust per animal. Apply on back and neck of the animals and repeat applications at 10-14 day intervals. Do not apply to Dairy Cattle later than 5 hours before milking or during milking. Do not contaminate water troughs.

POULTRY- (Chickens, Ducks, Geese and Turkeys): To control Poultry Lice, Northern Foul Mite, and Chicken Red Mite, dust individual birds by means of a shaker can or hand duster. Repeat in 4 to 8 weeks when necessary. Premise Treatment: Apply 1 pound per 50 square feet of litter and floor space and to nests, roost and adjacent areas.

HOGS- For lice control: Treat animal thoroughly and repeat in 10 days and thereafter as needed. Avoid contamination of feed, food containers, and watering troughs.

CAUTION

Do not treat animals under one month of age. Do not apply to lactating dairy animals or within 2 weeks of freshening. Do not contaminate feed, food containers, and watering troughs.

CATTLE- Lactating Dairy Cattle: Horn Fly, apply 3 tablespoons of 5% Malathion dust per animal. Apply on back and neck of the animals and repeat applications at 10-14 day intervals. Do not apply to Dairy Cattle later than 5 hours before milking or during milking. Do not contaminate water troughs.

POULTRY- (Chickens, Ducks, Geese and Turkeys): To control Poultry Lice, Northern Foul Mite, and Chicken Red Mite, dust individual birds by means of a shaker can or hand duster. Repeat in 4 to 8 weeks when necessary. Premise Treatment: Apply 1 pound per 50 square feet of litter and floor space and to nests, roost and adjacent areas.

HOGS- For lice control: Treat animal thoroughly and repeat in 10 days and thereafter as needed. Avoid contamination of feed, food containers, and watering troughs.

CAUTION

Avoid contact with skin. Avoid breathing dust. Wear a respirator of a type approved by the U.S. Bureau of Mines when handling this product. Harmful if swallowed. Wash thoroughly with soap and water after use.

NOTE TO PHYSICIAN: Malathion upon prolonged use, may cause cholinesterase inhibition. Atropine is antidotal.

Keep away from children and domestic animals. Avoid contamination of feed and foodstuffs.

Highly toxic to fish. Do not contaminate any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Avoid application to certain ferns, including Boston, Maiden Hair, and Pteris as well as some species of Crassula



411-22

EPA Est. 635-MI-1; OH-1

E.P.A. Reg. No. 635-430