

# E-Z FLO PARATHION 15-W

**DANGER:**  
KEEP OUT OF THE REACH OF CHILDREN



**POISON**



**POISON**

635-431  
6-9-70

**ACTIVE INGREDIENTS:**

Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate) ..... **15.0%**

**INERT INGREDIENTS:** ..... **85.0%**

**100.0%**

**DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.**

A wettable powder for use in water as an insecticide

## DIRECTIONS FOR USE

Application should be made only by experienced personnel using approved safety equipment. For recommended amount of this material on surface of water, in nearly filled spray tank or premixing bucket, half-filled with water and pour into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

**RECOMMENDATIONS:**

All recommendations are given in terms of pounds of this material per 100 gallons of water, unless stated otherwise. Thorough full coverage applications should be made.

**APPLE:** Codling Moth, Plum Curculio, Red-banded Leaf Roller, second and third brood, Pear and Woolly Aphids—1 1/4 lbs.; Bud Moth, Red Bug, Red-banded Leaf Roller, first brood, Rusy & Green Apple aphids and mites—1 lb.

Repeat applications at 7 to 10 day intervals may be necessary for the control of Codling Moth, McIntosh and related varieties of the Golden Delicious and Jonathan varieties. Full coverage spray and do not apply within 14 days of harvest.

**CHERRY:** Mites & Aphids—1 1/2 to 1 lb. Repeat applications at 7 to 10 day intervals may be necessary for Spider Mites.

Plum Curculio, Cherry Fruitworm, Oriental Fruit Moth, Bud Moth, Pearworm, Pear, Jade Scale & Pear Scale—1 1/2 lbs. For San Jose Scale, apply at time crawlers emerge. Use as full coverage spray and do not apply within 14 days of harvest.

**PEACH:** Cottony Peach Scale, Aphids & Spider Mites—1 1/2 to 2 lbs. Oriental Fruit Moth, Pear and facing insects & scales—1 1/2 to 2 lbs. Apply after all eggs have hatched for Pear and Scale.

Use as a full coverage spray and do not apply within 14 days of harvest. Do not apply more than 1 1/2 lbs. total parathion 15-W per acre per year. Do not apply in California.

**BEANS:** Mexican Bean Beetle, Leafhopper, Aphids, Red Spider Mite & Armyworm. Use in sufficient water for thorough coverage. Repeat application at 7 to 10 day intervals until controlled. Do not apply more than 1 1/2 lb. per acre per year. Do not apply within 14 days of harvest.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Diamond-Back Moth, Imported Cabbage worm and Armyworms—1 1/4 lbs. per acre. Aphids & Thrips—1 lb. per acre.

Use in sufficient water for thorough coverage. Repeat application at 7 to 10 day intervals until controlled. Do not apply more than 3 lb. per acre application.)

DO NOT APPLY WITHIN 7 DAYS OF HARVEST ON BROCCOLI, BRUSSELS SPROUTS AND CAULIFLOWER; 10 DAYS OF HARVEST ON CABBAGE. NOTE: IN FLORIDA, DO NOT APPLY ON ABOVE COLE CROPS WITHIN 14 DAYS OF HARVEST.

**CUCUMBER, SQUASH, MELON:** Cucumner Beetle, Melonworm, Pickleworm, Aphid, Stink Bug and Vine Borer, 2 lbs. per acre. Use in sufficient water for thorough coverage. Repeat application at 7 to 10-day intervals until controlled. Do not apply more than 3 lb. per acre application.)

DO NOT APPLY BEFORE PLANTS START TO VINE. DO NOT APPLY UNLESS PLANTS ARE DRY. DO NOT APPLY WITHIN 15 DAYS OF HARVEST ON CUCUMBERS AND SQUASH; 7 DAYS OF HARVEST ON MELONS.

**POTATO, TOMATO:** Colorado Potato Beetle, Flea Beetle, Leafhopper, Serpentine Leaf Miner, Grasshopper, Whitefly & Aphids (Apply with usual fungicide, 1 lb. per acre. Armyworms—1 1/4 lbs. per acre. Use in sufficient water for thorough coverage. Repeat application at 7 to 10 day intervals until controlled. Do not apply more than 3 lb. per acre application.)

DO NOT APPLY TO POTATO WITHIN 5 DAYS OF HARVEST OR TO TOMATO WITHIN 10 DAYS OF HARVEST.

**TOBACCO: Seedbeds:** Aphids—1 lb. per acre. Field Aphids & Suckfly—1 lb. per acre. Use in sufficient water for thorough coverage. Repeat application at 7 to 10 day intervals until controlled.

MAKE NO APPLICATION WITHIN 5 DAYS OF PRIMING OR 15 DAYS OF CUTTING. AVOID GETTING WET WITH PLANT JUICES WHEN CUTTING.

**DANGER**

# E-Z FLO PARATHION 15-W

6-9-70  
131

## ACTIVE INGREDIENTS:

Parathion (O,O-diethyl O-p-nitrophenyl  
thiophosphate)

15.0%

## INERT INGREDIENTS:

85.0%

100.0%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

## DIRECTIONS FOR USE

Read the label carefully before using this product. It contains important information on the safe and effective use of this product. Do not use this product if the label is missing or if the instructions are not followed. Do not use this product on crops or plants that are not listed on the label. Do not use this product on crops or plants that are listed on the label if the instructions are not followed. Do not use this product on crops or plants that are listed on the label if the instructions are not followed.

### RECOMMENDATIONS

- APPLE
- CHERRY
- PEACH
- BEANS