

ACCEPTED

APR 1 1971

U.S. DEPARTMENT OF AGRICULTURE
FEDERAL REGISTER
68 434
SUBJECT

NET CONTENTS

E-Z FLO Superkill Parathion 1 Thiodan® 2 EC

(Contains 1 Lb. Parathion and 2 Lbs. Thiodan® Per Gallon)



POISON



DANGER:

KEEP OUT OF THE REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNINGS
AND ANTIDOTE STATEMENTS

ACTIVE INGREDIENTS:

Parathion*	11.5%
Endosulfan**	22.8%
Xylene	60.0%
INERT INGREDIENTS:	5.7%
	100.0%

*O, O-Diethyl O-p-nitrophenyl thiophosphate
**Thiodan (Hexachlorocyclohexane 2, 4, 3-benzodioxathiepin oxide)
Thiodan is a registered trademark of Canadian Hoechst Limited.
U. S. Patent No. 2,799,685.

The manufacturer makes no warranties, express or implied, concerning this product for its use, which extend beyond the description on the label. All statements made concerning this product apply only when used as directed.

SOLD BY

E-Z FLO CHEMICAL COMPANY

DIVISION OF GROWER SERVICE CORPORATION

ABERDEEN, NORTH CAROLINA

COLUMBUS, OHIO

LANSING, MICHIGAN

072

53-263-804

USDA Reg No 635-434

DIRECTIONS FOR USE

**BEFORE USING STUDY SAFETY DIRECTIONS
DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE. BURN OR BURY
STAY OUT OF SMOKE OR FUMES**

Apply the dosages indicated per acre in 10 to 150 gals. of water through conventional ground equipment to give thorough coverage of the plants.

BROCCOLI: To control Cabbage Aphids, Cabbage Looper, Imported Cabbageworm, Diamond-back moth larvae, Cross striped Cabbage worm, Flea Beetle, Harlequin Bug, Stink Bug, use 1½ to 2 quarts per acre. Do not apply within 7 days of harvest.

CABBAGE: To control cabbage looper, imported cabbage worm, diamond-back moth larvae, cross striped cabbageworm, use 1 to 1½ quarts per acre. To control flea beetle, harlequin bug, stink bug use 1½ quarts per acre. NOTE: Do not apply within 14 days of harvest.

COTTON: To control spider mites, desert spider mite, tumid spider mite, two spotted spider mite, bollweevil, aphids, cotton leafworm, lygus bug, bollworm, cabbage looper--use 2 quarts per acre. NOTE: Do not apply after bolls begin to open nor within 5 days of hand picking. Do not graze dairy or meat animals in treated fields.

CUCUMBERS, PUMPKINS, SUMMER SQUASH, WINTER SQUASH: To control aphids, cucumber beetles use 2 quarts per acre. NOTE: Do not apply before vining or within 15 days of harvest on cucumbers, nor within 15 days of harvest on squash, nor within 10 days of harvest on pumpkins.

HEAD LETTUCE: To control cabbage looper, green peach aphid, diamond-back moth larvae, imported cabbage worm use 1½ to 2 quarts per acre. Do not make more than 3 applications after thinning. Do not apply within 14 days of harvest. Remove outside leaves at harvest. Do not feed crop waste to livestock.

MELONS: To control aphids, cucumber beetles use 1 to 2 quarts per acre. To control cabbage looper, omnivorous leaf roller use 2 quarts per acre. Do not apply within 7 days of harvest.

PEPPERS: To control flea beetle, green peach aphid use 1 quart per acre. NOTE: Do not apply within 15 days of harvest.

POTATOES: To control flea beetle, Colorado potato beetle, leafhopper, aphids, southern armyworm, green stink bug, leaf footed plant bugs and other plant bugs, potato tuberworm, whitefly, use 1 to 2 quarts per acre. To control false chinch bug use 2 quarts per acre. To control potato psyllid use 1½ to 2 quarts per acre. NOTE: Do not apply within 5 days of harvest.

TOBACCO: To control budworm, tobacco suckfly, cabbage looper, flea beetle, green peach aphid and hornworm use 1 quart per acre. NOTE: Do not apply within 5 days of priming or within 15 days of cutting. Do not enter fields within 24 hours of application. Protective clothing should be worn during application. Workers should avoid having plant juices coming in contact with their skin.

TOBACCO (plant bed treatment): Green June bug larvae--1 pint per 125 gals. At first sign of larvae activity apply as a drench using 1 gallon of spray mixture per 10 square yards of plant bed.

TOMATOES: To control aphids, blister beetle, Colorado potato beetle, flea beetle, green stink bug, whitefly use 1 quart per acre. NOTE: For hornworm--use 1 2 quarts/acre. Do not apply within 10 days of harvest.

REGISTERED

65-134

E-Z FLO Superkill Parathion 1 Thiodan 2 EC

(Contains 1 Lb Parathion and 2 Lbs Thiodan Per Gallon)

ACTIVE INGREDIENTS:	
Parathion*	11.5%
Endosulfan**	22.8%
Xylene	60.0%
INERT INGREDIENTS:	
	5.7%
	<hr/> 100.0%

SOLD BY
E-Z FLO CHEMICAL COMPANY
 DIVISION OF GROWER SERVICE CORPORATION
 GREENSBORO, NORTH CAROLINA COLUMBUS, OHIO LANSING, MICHIGAN

DIRECTIONS FOR USE

BEFORE USING STUDY SAFETY DIRECTIONS
 NEVER REUSE CONTAINER WHEN EMPTY NEVER RE USE BURN OR BURY
 STAY OUT OF SMOKE OR FUMES

BEANS: ...
BELL PEPPERS: ...
BROCCOLI: ...
CUCUMBERS: ...
EGGPLANTS: ...
KALE: ...
MELONS: ...
PUMPKINS: ...
SQUASH: ...
TOMATOES: ...