

E-Z FLO MSMA PLUS SURFACTANT

(CONTAINS 4 LBS. MSMA PER GALLON)

CAUTION:
KEEP OUT OF THE REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:
Monsodium Acid methanearsonate 35.42%
INERT INGREDIENTS: 64.58%
100.00%

Total arsenic, all in water-soluble form, expressed as elemental — 16.38%

Seller guarantees and warrants (a) that the active ingredient content and the total net weight are as stated within lawful limits and (b) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops, and upon reports of field experience. Tests have not been made on all varieties or in all states. Seller makes no other express or implied guarantee, warranty or representation. No Warranty of Merchantability or of Fitness for a Particular Purpose is implied.

SOLD BY
E-Z FLO CHEMICAL COMPANY
DIVISION OF GROWER SERVICE CORPORATION
ABERDEEN, NORTH CAROLINA COLUMBUS, OHIO LANSING, MICHIGAN

88-23

53-263-811

U.S.D.A. Reg. No. 635-479

DIRECTIONS FOR USE

HERBICIDE FOR POST EMERGENCE WEED CONTROL IN
COTTON AND NON-CROP AREAS

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use of this material in a manner or at any time other than in accordance with the directions may produce plant injury, excessive residues, or other undesirable results.

DIRECTIONS FOR USE

COTTON — MSMA is useful as a directed spray application to weeds in cotton for post-emergence control of weeds and grasses including Johnson grass, nut grass, water grass, goose grass, dallis grass and cocklebur.

Mix at a rate of 2 quarts in 40 gallons of water. Apply at a broadcast rate of 40 gallons of this solution per acre. This is a combination of surfactant (wetting agent) and herbicide. It is unnecessary to add any other surfactant to the spray solution. For band application, apply at a rate of one gallon of this solution for each 1 inch of band width to be treated on cotton grown in 40-inch rows. A second application may be required during humid growing conditions. Time the second application, if necessary, about 1 to 3 weeks after the first application.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOOM. KEEP OFF COTTON FOLIAGE. DO NOT APPLY AFTER FIRST BLOOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK OR GRAZE TREATED AREAS.

Slight buring and a reddish discoloration of the cotton leaf may occasionally occur following the recommended treatment, but the cotton plant will develop normally.

NON-CROP — MSMA is useful for controlling Johnson grass, nut grass, water grass, goose grass and other weeds on drainage ditch banks, right-of-ways, storage yards, and similar non-crop areas.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION OR BY ANIMALS, OR WATER USED FOR IRRIGATION PURPOSES.

NOTE

KEEP FROM FREEZING. SOLIDS MAY APPEAR IN SOLUTION UNDER COLD STORAGE CONDITIONS. HOLDING SOLUTION AT ROOM TEMPERATURE, 70° F, A SHORT TIME REDISSOLVES ALL SOLIDS AND SOLUTION IS ACCEPTABLE FOR USE.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or foodstuffs. Keep children and animals off treated areas until the material is washed into the soil.

General Information

Mix at a rate of 2 quarts in 40 gallons of water. No additional surfactant is necessary. Spray the unwanted vegetation thoroughly to the point of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs, retreat as required.

APPLICATION METHODS

MSMA must be thoroughly dissolved. Fill the tank about half full with water, and with the pump operating add the required amount of herbicide. Finish filling the tank with water and apply.

If preferred, MSMA can be premixed by dissolving the required amount in 3 to 5 gallons of water in an open head bucket. This premix should then be placed in the partially filled spray tank and the tank filled.

Do not store spray solution in tank for a prolonged period. Clean application equipment thoroughly after use by flushing with water.

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

ACCEPTED

NET CONTENTS 4 GALLONS